



2014 Forest Grouse Parts Collection Summary



New forest grouse wing collection barrel by Tye Bridge, Douglas County, Oregon. The Umpqua Watershed District Office placed out 7 new collection barrels in Douglas County in the fall of 2014, for a total of 13 collection barrels in the Roseburg District.

Photo by Umpqua Watershed District Office, ODFW

Upland Game Bird Program
Oregon Department of Fish and Wildlife
4034 Fairview Industrial Dr SE
Salem, OR 97302
Ph: 503-947-6322
E-mail: kelly.m.walton@state.or.us

INTRODUCTION

Since 1980, wings and tails of forest grouse have been collected from hunters in Wallowa County. In 1984, collections for forest grouse were expanded to other counties in northeastern Oregon and portions of southwestern Oregon. Since that time the effort has increased to nearly statewide participation.

In 2014, wings and tails of blue¹ (*Dendragapus spp*), ruffed (*Bonasa umbellus*), and spruce grouse (*Falci pennis canadensis*) were obtained from 24 of the 36 counties in Oregon (Table 1). A total of 1,107 wings and tails were examined at 2 forest grouse wing bees. The total number of wings and tails obtained in 2014 was up 30% from the previous year and up 15% from the recent 5 year average of 959. Statewide blue grouse submissions were up 9% and ruffed grouse wings were up 43% compared to 2013. Spruce grouse wings represent a small proportion of all grouse wing submissions with only 3 wings from Wallowa County in 2014, a decrease from the recent 5 year average of 7 wings. There is no open spruce grouse season in Oregon.

¹Dusky and Sooty grouse considered collectively as “blue” grouse.

USE OF INFORMATION

Data from wings can be used by biologists to better understand the distribution and timing of grouse harvest in their areas (Figures 2 & 4), the relative proportions of harvest among species, the sex and age structure of the population, and the chronology of breeding activity (Figures 3 & 5). In comparison to other methods for tracking trends in population size and productivity, such as brood route and hunter harvest telephone surveys, data gathered from wings is an effective and low cost method for monitoring trends (Hansen et al. 2015).

Distribution and timing of harvest have relevance to obtaining information about grouse populations, season structure, and to hunter participation for coordination of law enforcement activities. Sex and age data reveal the reproductive performance in a population (productivity), and in conjunction with abundance information, provide insight into population trends. Hatching data may be used to understand the timing of reproduction in specific areas and provide information to develop appropriate census procedures. For example, if hatching times

Table 1. Forest grouse wings submitted to the 2014 Oregon forest grouse wing bees at Ladd Marsh Wildlife Area on January 27, 2015 and the Umpqua Watershed District Office (Roseburg) on March 17, 2015. Wing bee counts from 2013 are provided for comparison.

County	"Blue" Grouse		Ruffed Grouse		Spruce Grouse		Total Wings		% change from 2013
	2014	2013	2014	2013	2014	2013	2014	2013	
Baker	34	21	17	12			51	33	54.5
Deschutes		1	1				1	1	0.0
Grant	47	38	122	124			169	162	4.3
Harney	1	14		6			1	20	-95.0
Jefferson		2					0	2	-100.0
Klamath	2	5		2			2	7	-71.4
Lake	40	26					40	26	53.8
Morrow	14	2	15	1			29	3	866.7
Umatilla	3	2		4			3	6	-50.0
Union	37	53	121	71			158	124	27.4
Wallowa	59	93	157	174	3	3	219	270	-18.9
Wheeler	1		1				2	0	
Clatsop	2		2	5			4	5	-20.0
Coos	4		11				15	0	
Curry	6	3	7	9			13	12	8.3
Douglas	75	25	233	64			308	89	246.1
Hood River	6	14	12	8			18	22	-18.2
Jackson			3	2			3	2	50.0
Lane	9	10	8	7			17	17	0.0
Lincoln			2				2	0	
Linn	6	1	2	1			8	2	300.0
Marion	7	11	3	5			10	16	-37.5
Tillamook	5	7	1	1			6	8	-25.0
Wasco	8	8	18	16			26	24	8.3
Washington				1			0	1	-100.0
Yamhill			2	2			2	2	0.0
Total	366	336	738	515	3	3	1,107	854	29.6

differed substantially among regions of the state, the timing of summer censuses could be adjusted because the probability of observing a bird is a function of bird age and habitat conditions. Samples obtained through hunter harvest may not adequately reflect sex and age ratios of a population and may change through the course of the season. Age ratios during the first two weeks of the season provide the best index to reproduction, while there is no

significant change in sex ratios during the course of the season (Hansen et al. 2012). This report provides age ratios for the first two weeks of the season and for the entire season. The age ratios for the entire season will allow comparison to data collected in previous years.

METHODS

District wildlife biologists collect grouse parts from hunters by placing “wing barrels” in locations where grouse hunters are likely to encounter them. Bags are placed at the barrels which instruct hunters to remove one wing and the tail from each grouse they kill and place it in a single bag. They are also asked to record the date, county and general location of the kill. Barrels are checked periodically throughout the season and any bags not dated or labeled by hunters are labeled with the barrel location and date of collection. This year, the Umpqua Watershed District Office (Roseburg) increased collection effort by placing 7 new forest grouse wing collection barrels out in Douglas County for a total of 13 collection barrels in that district.

Field staff also distributed wing bags to known grouse hunters. Additionally, wing bags are mailed to a list of cooperating hunters by wildlife division staff prior the hunting season and an advertisement requesting participation in the program is placed in the annual game bird regulations and on the department’s website. Finally, further opportunities to solicit participation such as magazine articles and newspaper interviews are utilized when available.

Each winter biologists gather at wing bees to collect information from the parts. In 2014-15, wing bees were held at Ladd Marsh Wildlife Area (Figure 1 &2) and the Umpqua Watershed District Office. Data collected from each set of parts are: species, location of kill, date of kill, sex, age, and the stage of primary wing feather molt for immature birds. Age is recorded as adult or immature and in addition, the adult class is further subdivided to adult or yearling (if discernable). After the wing bees, data is entered into a spreadsheet which contains formulas for estimating the age, in days, of immatures based on the sequential replacement of primary wing feathers. Hatch dates are then back calculated for birds of known harvest date; provided they were harvested on or prior to 10 October (immature molt of primaries 1 through 8 is usually complete by 10 October).



Figure 1. Determining the age and sex of grouse parts received at the eastside wing bee at Ladd Marsh Wildlife Area, 27 January 2015 (Martyne Reesman, Kelly Walton, Mike Moore, Chase Brown).



Figure 2. Phillip Perrine showing off an immature female “blue” grouse wing at the eastside wing bee at Ladd Marsh Wildlife Area, 27 January 2015.

BLUE GROUSE RESULTS

During 2014, 366 wings and tails from “blue” grouse were collected in Oregon, an increase of about 9% from the previous year and a 16% decline from the recent 5-year average of 436 wings. The 2014 hunting season allowed a daily bag limit of 3 birds with 9 in possession. The season started 1 September statewide and ended 31 December in eastern Oregon and ended 31 January in western Oregon, which includes Hood River and Wasco counties. Thirty percent of the wings and tails were harvested during the first week of the season and 57% were harvested by the end of September. Similar to most previous seasons, there was a second peak of submissions during late September and early October; the beginning of many firearm deer and elk seasons (Figure 3).

Sixty-five percent (238) of the wings and tails were obtained from eastern Oregon. The majority of submissions from eastern Oregon were from Wallowa (25%), Grant (20%), and Lake (17%) counties. The remainder of the eastern Oregon submissions came from 7 other counties. One hundred twenty eight wings and tails were obtained from 10 counties in western Oregon, a 62% increase from the number received in the previous year. The majority of the submissions were from Douglas (56%) county.




Figure 3. The number of sooty and dusky grouse wing/ tail collections, by week of reported harvest during 2014-15 Oregon hunting season.

Age and Sex Ratios

Immature grouse comprised 79% of the wings for “blue” grouse during the first two weeks of the season and 72% of the sample from the entire season (Table 2). The proportion of immatures in the harvest indicates above average (~65% immature) production. Males represented 43% of the statewide sample, 56% of adult sample, and 38% of immatures in 2014.

Hatching Chronology

Statewide, hatch dates for dusky and sooty grouse harvested during the 2014 hunting season ranged from 5 May to 15 July, which is similar to previous years. Dusky grouse hatch dates ranged from 5 May to 15 July (\bar{x} = 2 June) and sooty grouse hatch dates ranged from 10 May to 13 July (\bar{x} = 5 June). For dusky grouse, 75% hatched between 18 May and 16 June, while 75% of sooty grouse hatched between 22 May and 17 June. Typical of most years, the peak sooty grouse hatch was later than the peak dusky grouse hatch. Mean hatch dates were

average (Figure 4) and were earlier than hatch dates in 2010 and 2011, which were among the latest recorded during this study (1980 – 2014).




Figure 4. Week of hatch for dusky and sooty grouse in Oregon as estimated from primary feathers from hunter-harvested wings during 2014. Hatch dates were only estimated for birds that were harvested on or before October 10.

Table 2. “Blue” grouse sex ratios, age ratios and hatch dates by species in 2014 from hunter submitted wings. Ratios are presented for the first two weeks of season and for the entire season. Wings and tails harvested during the first two weeks are the best indicator for age ratios (Hansen et al. 2012). Hatch date is only estimated for grouse harvested on or before October 10.

	<u>Gender Ratios</u>				<u>Age Ratios</u>			<u>Hatch Dates</u>	
	n	M:F	AM:AF	IM:IF	n	I:A	I:AF	n	Mean, Range
Dusky									
1st 2 wks	80	46:54	71:29	41:59	81	83:17	94:6	---	---
Total	193	49:51	67:33	72:28	194	74:26	89:11	88	June 2, May 5 - July 15
Sooty									
1st 2 wks	54	24:76	21:79	25:75	54	74:26	78:22	---	---
Total	159	37:63	45:55	34:66	165	71:29	82:18	63	June 5, May 10 - July 13
All									
1st 2 wks	134	37:63	46:54	35:65	135	79:21	88:12	---	---
All Total	352	43:57	56:44	38:62	359	72:28	86:14	151	June 4, May 5 - July 15

Wallowa County – 1980 to 2014

From 1980 through 2014 hunters in Wallowa County submitted parts representing 9,165 dusky grouse. In 2014, 59 wings were collected, which is down 78% from the average since the wing bee began in 1980. Males comprised 78% of the sample, which is higher than the long-term average of 66% (Table 3).

The immature proportion (69%) of the harvest was above average and suggests good production as the proportion of immatures was above 65% (Table 3). Production has been above 65% for the past two years (2013 and 2014) after a period of lower production (2010-2012). The proportion of immatures in Wallowa County has ranged from a low of 38% in 1982 to a high of 76% in 2008 (1980–2014, \bar{x} = 61%). Over the past 35 years, dusky grouse populations in Wallowa County appeared to have 4 years of poor production as measured by a proportion of immatures <50% (1982, 1991, 1995, and 2006) and 13 years of good production as measured by a proportion of immatures >65% (1980, 1983, 1985, 1986, 1989, 1996, 1998, 2000, 2007, 2008, 2009, 2013, 2014; Table 3).

Mean hatch dates have ranged from 25 May (1992, 1998, and 2004) to 6 June (1995 and 1999) with the preponderance of young hatched during a 3-week interval between late May

and early June (Table 3). In 2014, the mean hatch date was 3 June and ranged from 13 May to 24 June.

Table 3. Sex ratios, age ratios and hatching dates of dusky grouse determined from parts submitted by hunters from harvest in Wallowa County, Oregon, 1980 to 2014.

Season	<i>n</i>	Sex Ratios			Age Ratios		Hatch Information			
		M:F	AM:AF	IM:IF	I:A	I:AF	Mean	Range		
1980	59	54:46	83:17	41:59	69:31	93:7				
1981	125	57:43	60:40	55:45	62:38	80:20	29-May	7-May	to	29-Jun
1982	95	53:47	53:47	53:47	38:62	56:44	31-May	16-May	to	16-Jun
1983	165	53:47	57:43	51:49	72:28	86:14	30-May	8-May	to	25-Jun
1984	155	57:43	63:37	53:47	52:48	74:26	4-Jun	13-May	to	8-Jul
1985	258	53:47	63:37	49:51	72:28	88:12	1-Jun	4-May	to	4-Jul
1986	598	58:42	74:26	52:48	70:30	90:10	26-May	3-May	to	15-Jul
1987	736	58:42	72:28	51:49	65:35	87:13	26-May	2-May	to	14-Jul
1988	471	54:46	60:40	47:53	53:47	75:25	2-Jun	28-Apr	to	19-Jul
1989	371	53:47	59:41	51:49	70:30	85:15	30-May	29-Apr	to	10-Jul
1990	286	58:42	65:35	55:45	54:46	77:23	27-May	5-May	to	1-Jul
1991	260	60:40	68:32	50:50	43:57	70:30	1-Jun	9-May	to	13-Jul
1992	284	54:46	61:39	47:53	57:43	78:22	25-May	2-May	to	26-Jun
1993	200	58:42	61:39	57:43	65:35	83:17	2-Jun	10-May	to	28-Jun
1994	249	59:41	66:34	52:48	58:42	80:20	28-May	10-May	to	21-Jun
1995	140	47:53	61:39	30:70	43:57	66:34	6-Jun	14-May	to	10-Jul
1996	261	61:39	75:25	54:46	67:33	89:11	30-May	10-May	to	8-Jul
1997	205	54:46	78:22	41:59	61:39	88:12	30-May	10-May	to	24-Jun
1998	361	59:41	73:27	53:47	66:34	88:12	25-May	8-May	to	30-Jun
1999	453	59:41	69:31	51:49	59:41	82:18	6-Jun	11-May	to	5-Jul
2000	379	60:40	82:18	51:49	68:32	92:8	27-May	3-May	to	3-Jul
2001	570	52:48	62:38	47:53	65:35	83:17	31-May	3-May	to	7-Jul
2002	376	59:41	64:36	56:44	63:37	83:17	5-Jun	5-May	to	29-Jul
2003	460	64:36	74:26	58:42	65:35	88:12	3-Jun	6-May	to	17-Jul
2004	251	50:50	56:44	47:53	51:49	70:30	25-May	5-May	to	30-Jun
2005	209	64:36	80:20	56:44	59:41	88:12	1-Jun	9-May	to	14-Jul
2006	163	61:39	70:30	54:46	48:52	76:24	1-Jun	10-May	to	8-Jul
2007	172	55:45	55:45	56:44	70:30	84:16	27-May	6-May	to	4-Jul
2008	104	53:47	56:44	53:47	76:24	88:12	5-Jun	10-May	to	22-Jul
2009	173	58:42	64:36	55:45	68:32	87:13	30-May	9-May	to	12-Jul
2010	128	47:53	58:42	38:62	55:45	76:24	5-Jun	9-May	to	6-Jul
2011	150	57:43	61:39	46:54	57:43	83:17	5-Jun	8-May	to	15-Jul
2012	126	46:54	66:34	29:71	52:48	76:24	1-Jun	15-May	to	26-Jun
2013	93	61:39	65:35	59:41	66:34	85:15	2-Jun	5-May	to	30-Jun
2014	59	44:56	78:22	46:54	69:31	91:9	3-Jun	13-May	to	24-Jun

RUFFED GROUSE RESULTS

In 2014, a total of 738 ruffed grouse wings and tails were collected in Oregon, a 43% increase from 515 wings last year and a 64% increase from the recent 5-year average of 451 wings. The 2014 ruffed grouse hunting season allowed a daily bag limit of 3 birds with 9 in possession. Statewide the season began 1 September and extended through 31 December in eastern Oregon and through 31 January 2015 in western Oregon, which includes Hood River and Wasco counties. Typically a large portion of the ruffed grouse wings are submitted during the first couple weeks of season, with a second peak in harvest occurring during the start of many firearm big game seasons, usually the first week in October. In 2014, the peak of ruffed grouse wing returns occurred during the second week of the grouse season and again during the start of the big game rifle seasons (Figure 4). Eleven percent of the ruffed grouse parts were submitted during the first week of the season. In prior years the number of grouse wings returned dropped considerably after the first week of big game rifle seasons. In 2014, a large percentage of the wings were collected in October, with 40% of wings harvested in September and 42% in October (Figure 5).

In eastern Oregon, 434 samples were collected, a 10% increase from 2013 and above the recent 5-year average of 383. Most of the samples collected in eastern Oregon were from Grant, Union, and Wallowa Counties. In western Oregon, 304 ruffed grouse samples were collected, a 151% increase over last year. Additional wing collection effort in Douglas County accounted for the majority of the increase. In 2014, 233 ruffed grouse wings were collected in Douglas County compared to the recent five year average of 55 wings. Wings were collected from a total of 12 counties in western Oregon.

Age and Sex Ratios

Because of the lack of tails, or the rump feathers attached to the tail, gender could not be determined for 45% of the ruffed grouse submissions. Age was determined for 98% of the samples obtained from eastern and western Oregon.




Figure 5. Timing of Oregon’s 2014 ruffed grouse harvest as estimated from 697 ruffed grouse wings with harvest dates.

Immature grouse comprised 74% of the statewide sample during the first two weeks of the season indicating good production. Most (86%) of the early season ruffed grouse wings were collected in eastern Oregon. Age ratios during the first two weeks of the season provide the best index to reproduction because age ratios in the harvest can change over the course of the season (Hansen et al. 2012). For the entire season, immatures comprised 56% of the statewide harvest of ruffed grouse. In eastern Oregon, 68% of ruffed grouse samples were from immatures and 39% of submissions from western Oregon were immatures (entire season; Table 4). The proportion of immatures in western Oregon indicates below average production. An analysis of wings from the previous 27 years in Oregon found that immatures accounted for 33–74% of the sample. Ruffed grouse populations in other states have also displayed highly variable productivity that ranged from 39–80% of immatures in fall populations (Dorney 1963, Davis and Stoll 1973). The differences in production may be related to local variations and naturally occurring population cycles.

Males of all ages accounted for 48% of the wings collected. The proportion of adult wings that were male was 50% and 47% for eastern and western Oregon, respectively. Males

accounted for 55–61% of the adult population in several states in the mid-West (Dorney 1963, Davis and Stoll 1973, Major and Olson 1980). In 2014, males were 45% of the immature birds submitted from eastern Oregon. Confidence in the sex ratios would be improved if sex could be determined for a higher proportion of the samples. In 2014, 55% of the ruffed grouse samples included diagnostic feathers for sex identification.

Hatching Chronology

The mean hatch date for ruffed grouse collected during the 2014 hunting season was 2 June. Hatch dates were estimated from 224 wings, of which 24 were from western Oregon. Given the small sample from the west side, wings from both sides of the state were pooled for analysis (Table 4). Similar to “blue” grouse, peak hatch dates for ruffed grouse were close to average. Hatching dates ranged from 3 May to 22 July (\bar{x} = 2 June), with 75% ruffed grouse in the harvest hatching between 21 May and 14 June (Figure 6).




Figure 6. Timing of Oregon’s 2014 ruffed grouse hatch as estimated from 224 wings harvested on or before 10 October.

Table 4. Sex ratios, age ratios and hatching dates of ruffed grouse as determined from parts submitted by hunters from Oregon harvest during the 2014-15 hunting season.

	<u>Gender Ratios</u>			<u>Age Ratios</u>		<u>Hatch Dates</u>	
	n	M:F	AM:AF	IM:IF	I:A	I:AF	n Mean, Range
East	199	50:50	58:42	45:55	68:32	92:8	200 June 2, May 3 - July 22
East, 1st 2 wks	75	51:49	53:47	50:50	79:21	94:6	--- ---
West	209	47:53	50:50	42:58	39:61	63:37	24 June 4, May 10 - July 9
Statewide	408	48:52	53:47	44:66	56:44	81:19	224 June 2, May 3 - July 22

Wallowa County - 1983 to 2014

From 1983 through 2014, hunters have submitted 6,198 ruffed grouse wings and tails from Wallowa County. The proportion of immatures (76%) in the 2014 wing sample was up from last year (61%) and was the highest since 1983. Wing data collected since 1983 in Wallowa County indicated exceptional production from 1983–1990, then age ratios declined and stabilized until they again increased in 1999 through 2001. Age ratios have since been stable and slightly higher than during the early 90s except for 2004 and 2010, the age ratio in 2014 indicates exceptional production. The ratio of males (62%) in the sample was higher than last year (48%) and above the long-term average (~57%), however only half the submissions contained the diagnostic feathers to determine sex. A large proportion of hunter submissions continue to lack the diagnostic rump feathers or other keys to gender identification. The mean hatch date of 30 May was similar to 2013 and similar to the long term mean.

Table 5. Sex ratios, age ratios and hatching dates of ruffed grouse as determined from parts submitted by hunters from grouse harvested in Wallowa County, Oregon, 1983 to 2014.

Season	n	Sex Ratios			Age Ratios		Hatch Dates		
		M:F	AM:AF	IM:IF	I:A	I:AF	Mean	Range	
1983	70				83:17				
1984	47	50:50	0:100		66:34	97:3	5-Jun	22-May	to 20-Jun
1985	193	56:44	64:36	52:48	75:25	92:8	28-May	3-May	to 7-Jul
1986	395	61:39	69:31	56:44	72:28	93:7	29-May	5-May	to 14-Jul
1987	372	59:41	51:49	64:36	70:30	88:12	27-May	4-May	to 28-Jun
1988	212	69:31	78:22	64:36	68:32	95:5	1-Jun	13-May	to 1-Jul
1989	139	55:45	50:50	57:43	74:26	90:10	2-Jun	2-May	to 29-Jun
1990	189	61:39	71:29	56:44	67:33	93:7	28-May	11-May	to 20-Jun
1991	155	64:36	62:38	65:35	63:37	88:12	3-Jun	7-May	to 6-Jul
1992	220	65:35	64:36	66:34	61:39	87:13	27-May	30-Apr	to 5-Jul
1993	55	65:35	71:29	60:40	62:38	86:14	1-Jun	15-May	to 2-Jul
1994	112	53:47	52:48	54:46	55:45	76:24	25-May	12-May	to 26-Jun
1995	84	61:39	68:32	52:48	57:43	89:11	30-May	12-May	to 26-Jun
1996	180	62:38	70:30	54:46	57:43	85:15	29-May	3-May	to 20-Jun
1997	169	61:39	84:16	34:66	58:42	92:8	31-May	3-May	to 18-Jun
1998	279	53:47	59:41	48:52	55:45	81:19	25-May	7-May	to 26-Jun
1999	370	44:56	48:52	41:59	64:36	89:11	2-Jun	8-May	to 6-Jul
2000	339	61:39	67:33	55:45	58:42	89:11	26-May	3-May	to 21-Jul
2001	434	61:39	75:25	50:50	62:38	92:8	31-May	7-May	to 14-Jul
2002	165	51:49	60:40	42:58	56:44	83:17	5-Jun	11-May	to 7-Jul
2003	284	65:35	66:34	64:36	54:46	87:13	1-Jun	8-May	to 3-Jul
2004	98	48:52	57:43	35:65	49:51	76:24	28-May	7-May	to 18-Jun
2005	180	53:47	68:32	41:59	58:42	89:11	1-Jun	6-May	to 1-Jul
2006	152	56:44	62:38	48:52	59:41	87:13	26-May	5-May	to 10-Jul
2007	198	49:51	55:45	41:59	58:42	83:17	25-May	2-May	to 15-Jun
2008	94	56:44	61:39	52:48	63:37	87:13	4-Jun	7-May	to 27-Jun
2009	222	66:44	75:25	58:42	69:31	94:6	30-May	6-May	to 6-Jul
2010	167	54:46	56:44	50:50	43:57	73:27	6-Jun	14-May	to 28-Jun
2011	150	57:43	61:39	46:54	57:43	83:17	5-Jun	8-May	to 15-Jul
2012	143	47:53	51:49	41:59	55:45	68:32	30-May	6-May	to 1-Jul
2013	174	48:52	54:46	43:57	61:39	70:30	29-May	4-May	to 7-Jul
2014	157	62:38	64:36	61:39	76:24	93:7	30-May	3-May	to 29-Jun

SPRUCE GROUSE

Wing Collections – 1985 to 2014

Incidental to the harvest of dusky and ruffed grouse in Baker, Wallowa and Union counties, 193 spruce grouse wings and tails were collected from wing barrels from 1985 through 2014. Similar to last year, during 2014 3 spruce grouse wings were collected from Wallowa County. Wallowa County typically has the highest incidental harvest of spruce grouse but Union County has received more spruce grouse wings in recent years. During 1997, spruce grouse wings were obtained from Baker County for the first time, likely related to an increased effort in wing collection rather than range expansion. The 1988 wing bee recorded the highest number of spruce grouse wings (27). During the past 26 years, immatures and adults composed nearly equal proportions of the sample. In 2014, 3 immature wings were received. Oregon is on the southwest periphery of the natural range of spruce grouse and they are currently listed as vulnerable on Oregon's Sensitive Species List.

SUMMARY

Estimates of the statewide harvest from random phone surveys of upland game bird validation holders indicates harvest of "blue" grouse was similar to the previous year and estimates for ruffed grouse indicate a decrease in harvest from 2013 by 18%. Just over half of the ruffed grouse wings submitted in 2014 came from NE Oregon, estimated harvest for ruffed grouse for region 5 (Baker, Union, and Wallowa counties) decreased by 32% from 2013. Estimated dusky grouse harvest for region 5 increased by 13% compared to harvest the prior year.

For ruffed, sooty, and dusky grouse, mean hatch dates in 2014 were similar to 2013. Statewide, the proportion of immature "blue" grouse was higher (72%) than in 2013 (66%). The high proportion (>65%) of immature "blue" grouse suggests 2014 was another good year for recruitment. The proportion of immature ruffed grouse was similar (56%) to 2013 (57%) suggesting moderately good production.

2014 had the highest submission rate for wings in western Oregon since wing collections were initiated in 2000. The majority of the increase was a result of the increased collection

effort from 7 new wing barrels in Douglas County by the Umpqua Watershed. Sample submissions in the remainder of for western Oregon continue to remain disproportionately low. Even with increased collection efforts in western Oregon in 2014, about 39% of the submitted wings came from western Oregon, yet western Oregon accounted for 52% of the statewide “blue” grouse harvest and 66% of ruffed grouse harvest. For western Oregon, estimated harvest was down by 11% and 12% for “blue” and ruffed grouse, respectively. This underscores the need to continue to explore methods for increasing the submission rates for grouse harvested in western Oregon.

Statewide education efforts should continue to increase hunter awareness and participation for the need and value of returning wings and tails. These efforts should emphasize the need for hunters to include both a wing and a tail fan from “blue” grouse and a wing and a tail fan with attached rump feathers for ruffed grouse gender identification. Continuing hunter education efforts are critical for the success of future wing bees. Despite the desire for additional wings from Oregon forest grouse hunters, these wing collections still provide a valuable and reasonably low cost method of obtaining demographic profiles of grouse populations.

ACKNOWLEDGEMENTS

These data would simply not be available without the continued support and cooperation of Oregon hunters – for this we thank all the hunters who provided wings and tails! Forest grouse wing collection can also be a large workload for the wildlife districts, and their effort is greatly appreciated. Some wildlife districts have embraced this challenge and the results are self-evident – most of the forest grouse wings collected each year come from four NE Oregon counties and Douglas County– thank you.

LITERATURE CITED

- Davis, J.A., and R.J. Stoll, Jr. 1973. Ruffed grouse sex and age ratios in Ohio. *Journal of Wildlife Management*. 37:133-141.
- Dorney, R.S. 1963. Sex and age structure of Wisconsin ruffed grouse populations. *Journal of Wildlife Management*. 27:599-603.

Hansen, M. C., C. A. Hagen, D. A. Budeau, V.L. Coggins, and B.S. Reishus. 2015. Comparison of 3 surveys for estimating forest grouse population trends. *Wildlife Society Bulletin* 39:197–202

Hansen, M.C., C.A. Hagen, T.M. Loughin, D.A. Budeau, V.C. Coggins, and B.S. Reishus. 2012. Temporal Changes in Age and Sex Ratios of Forest Grouse Harvested in Northeast Oregon. *Journal of Wildlife Management*. 76:356-362.

Major, P.D., and J.C. Olson. 1980. Harvest statistics from Indiana ruffed grouse season. *Wildlife Society Bulletin*. 8:18-23.

Zwickel, F.C., J.H. Brigham, and I.O. Buss. 1975. Autumn structure of blue grouse populations in north-central Washington. *Journal of Wildlife Management*. 39:461-467