

Oregon Greater Sage-Grouse Population Monitoring: 2025 Annual Report




November 2025

Oregon Department of Fish and Wildlife

Skyler Vold

P.O. Box 8, Hines, OR 97738

Contact: Skyler Vold, 541-573-6582, Skyler.T.Vold@odfw.oregon.gov



Photos: Cover Page by Keith Kohl, retired ODFW, and Author Page by Skyler Vold, ODFW.

TABLE OF CONTENTS

PAGE

EXECUTIVE SUMMARY	1
OVERVIEW AND SPRING POPULATION MONITORING METHODS	1
Background on Lek Monitoring Efforts	1
Lek Monitoring Methods	2
Lek Data Analyses.....	6
LEK MONITORING EFFORT AND POPULATION ESTIMATES	8
Statewide.....	8
Abundance Results (2025), PAC-Scale	11
SUMMARY AND CONCLUSIONS	37
References	39
APPENDIX I – LONG-TERM POPULATION TRENDS BY PAC	41
Baker PAC.....	44
Beatys PAC	45
Brothers PAC	46
Bully Creek PAC.....	47
Burns PAC.....	48
Cow Lakes PAC	49
Cow Valley PAC.....	50
Crowley PAC.....	51
Diablo PAC	52
Drewsey PAC.....	53
Dry Valley – Jack Mountain PAC.....	54

Juniper Mountain PAC	55
Louse Canyon PAC.....	56
North Wagontire PAC.....	57
Paulina PAC	58
Pueblos PAC	59
Saddle Butte PAC	60
Sheepshead PAC	61
Soldier Creek PAC.....	62
Steens PAC	63
Trout Creeks PAC	64
Tucker Hill PAC	65
Warner PAC.....	66
Areas outside of PACs	67
APPENDIX II – 2025 ODFW ADOPT-A-LEK PROGRAM REPORT	68
Summary.....	68
2025 Volunteer Statistics	70
Gallery of Images	71
APPENDIX III – 2025 AERIAL SURVEY SUMMARY.....	75
Aerial Lek Searches Overview.....	75
2025 Aerial Survey	75
2024 Aerial Survey	75
2023 Aerial Survey	76
2022 Aerial Survey	76
2021 Aerial Survey	76

APPENDIX IV – 2025 SAGE-GROUSE WING-BEE REPORT (2024 HUNTING SEASON DATA)	78
Executive Summary	78
Overview	78
Age and Sex Composition	83
Nest Success and Production	88
Hatching Chronology	93
Annual Turnover	98
Conclusions	100
References	102

LIST OF TABLES

	<u>PAGE</u>
TABLE 1. DEFINITIONS OF ODFW’S SAGE-GROUSE LEK CONSERVATION STATUSES, 2025.....	5
TABLE 2. ESTIMATED SAGE-GROUSE ABUNDANCE (2025) AND PERCENT OF THE 2025 SPRING BREEDING SAGE-GROUSE POPULATION WITHIN EACH OF OREGON’S PRIORITY AREAS FOR CONSERVATION (PACS).....	12
<u>APPENDIX 1.</u>	
TABLE A1. 1. SAGE-GROUSE POPULATION TRENDS WITHIN EACH OF OREGON’S SAGE-GROUSE PRIORITY AREAS FOR CONSERVATION (PACS), 1980–2024.....	43
<u>APPENDIX 4.</u>	
TABLE A4. 1. SAGE-GROUSE HUNTING SEASON DATES, LENGTHS, DAILY AND SEASON BAG LIMITS, AND PERMITS AUTHORIZED, OREGON, 1993–2024.....	80
TABLE A4. 2. OREGON WILDLIFE MANAGEMENT UNITS OR SUB-UNITS WITH PERMITTED SAGE-GROUSE HARVEST, 2024.....	80
TABLE A4. 3. SEX COMPOSITION BY AGE CLASS, AND AGE COMPOSITION OF HARVESTED SAGE-GROUSE, ALL WILDLIFE MANAGEMENT UNITS OPEN TO HARVEST, OREGON, 1993–2024.	85
TABLE A4. 4. SEX COMPOSITION BY AGE AND AGE COMPOSITION OF HARVESTED SAGE-GROUSE WITHIN WILDLIFE MANAGEMENT UNITS WITH POTENTIAL SAGE-GROUSE HARVEST, OREGON, 2024. TOTAL BY PERCENT IN EACH AGE CLASS REPRESENTS THE OVERALL PROPORTION OF THE GIVEN AGE CLASS IN THE HARVEST.....	86
TABLE A4. 5. SAGE-GROUSE PRODUCTION DATA AS DETERMINED FROM HUNTER-HARVESTED WINGS, OREGON, 1982–1984, 1989–1992, 1993–2023, AND 2024. DATA IS DIVIDED INTO THESE PERIODS DUE TO THE SAGE-GROUSE HARVEST CLOSURE DURING 1985–1988, AND IMPLEMENTATION OF REVISED WING ANALYSIS METHODOLOGY IN 1993.....	87
TABLE A4. 6. SAGE-GROUSE NESTING SUCCESS AS INDICATED BY RETENTION OF AT LEAST PRIMARY FEATHER P9, ALL WILDLIFE MANAGEMENT UNITS WITH SAGE-GROUSE HARVEST, OREGON, 1993–2024.....	90
TABLE A4. 7. SAGE-GROUSE NEST SUCCESS AS INDICATED BY RETENTION OF AT LEAST PRIMARY FEATHER P9, AND PRODUCTION RATES, ALL WMUS WITH SAGE-GROUSE HARVEST, OREGON, 2024.....	91
TABLE A4. 8. ESTIMATED HATCH DATES FOR JUVENILE SAGE-GROUSE (% OF TOTAL) FROM HUNTER-HARVESTED WINGS, OREGON, 1993–2024.....	94
TABLE A4. 9. ESTIMATED HATCH DATES, FROM HUNTER-HARVESTED WINGS, FOR JUVENILE SAGE-GROUSE IN OREGON, 2024.....	95
TABLE A4. 10. ESTIMATED ANNUAL TURNOVER (%) OF ADULT SAGE-GROUSE, ASSUMING POPULATION STABILITY, OREGON, 1993–2024.....	99
TABLE A4. 11. ESTIMATED FALL SAGE-GROUSE POPULATION, MAXIMUM ALLOWABLE HARVEST, HUNTER STATISTICS, EXPECTED HARVEST, AND PERMIT ALLOCATION IN OREGON WILDLIFE MANAGEMENT UNITS (WMUS) WHERE SAGE-GROUSE HARVEST IS PERMITTED, 2024.....	102

LIST OF FIGURES

	<u>PAGE</u>
FIGURE 1. OREGON SAGE-GROUSE PRIORITY AREAS FOR CONSERVATION (PACS), 2025.	3
FIGURE 2. THE EFFECT OF TIME SINCE SUNRISE (PANEL A) AND DATE (PANEL B) ON THE PROBABILITY OF DETECTING MALE SAGE-GROUSE ATTENDING LEKS IN EASTERN OREGON, 2017–2025.	7
FIGURE 3. NUMBER OF SAGE-GROUSE LEKS AND LEK COMPLEXES KNOWN TO EXIST IN OREGON SINCE 1980, AND THE NUMBER OF LEKS AND LEK COMPLEXES SURVEYED EACH YEAR, 1980– 2025.	8
FIGURE 4. NUMBER OF SAGE-GROUSE LEK SURVEYS CONDUCTED PER YEAR IN OREGON (BLUE LINE), AND THE AVERAGE NUMBER OF SURVEYS CONDUCTED AT EACH SURVEYED LEK BY YEAR (ORANGE LINE), 1980–2025.	9
FIGURE 5. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN THE STATE OF OREGON, 2017–2025.	10
FIGURE 6. OREGON STATEWIDE SAGE-GROUSE POPULATION ESTIMATES, 1980–2025. THE SOLID BLACK LINE REPRESENTS THE ABUNDANCE ESTIMATES FROM THE NEW N-MIXTURE MODEL, AND THE DASHED GRAY LINE REPRESENTS THE POPULATION ESTIMATES FROM ODFW’S PREVIOUS MODEL. THE POPULATION GOAL IS SET AT THE 2003 LEVEL, REPRESENTING THE BASELINE FOR OREGON’S STATEWIDE SAGE-GROUSE POPULATION SIZE.	11
FIGURE 7. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN BAKER PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	13
FIGURE 8. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN BEATYS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	14
FIGURE 9. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN BROTHERS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	15
FIGURE 10. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN BULLY CREEK PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	16
FIGURE 11. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN BURNS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	17
FIGURE 12. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN COW LAKES PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	18
FIGURE 13. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN COW VALLEY PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	19
FIGURE 14. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN CROWLEY PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	20
FIGURE 15. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN DIABLO PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	21
FIGURE 16. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN DREWSEY PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	22
FIGURE 17. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN DRY VALLEY – JACK MOUNTAIN PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017– 2025.	23
FIGURE 18. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN JUNIPER MOUNTAIN PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	24
FIGURE 19. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN LOUSE CANYON PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	25
FIGURE 20. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN NORTH WAGONTIRE PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	26
FIGURE 21. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN PAULINA PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.	27

FIGURE 22. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN PUEBLOS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....28

FIGURE 23. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN SADDLE BUTTE PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....29

FIGURE 24. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN SHEEPSHEAD PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....30

FIGURE 25. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN SOLDIER CREEK PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....31

FIGURE 26. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN STEENS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....32

FIGURE 27. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN TROUT CREEKS PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....33

FIGURE 28. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN TUCKER HILL PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....34

FIGURE 29. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN WARNER PRIORITY AREA FOR CONSERVATION IN THE STATE OF OREGON, 2017–2025.....35

FIGURE 30. N-MIXTURE MODEL ESTIMATES OF SAGE-GROUSE ABUNDANCE WITHIN LOW-DENSITY HABITAT AREAS IN THE STATE OF OREGON, 2017–2025.....36

APPENDIX 1.

FIGURE A1. 1. SAGE-GROUSE POPULATION TRENDS IN THE BAKER PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....44

FIGURE A1. 2. SAGE-GROUSE POPULATION TRENDS IN THE BEATYS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....45

FIGURE A1. 3. SAGE-GROUSE POPULATION TRENDS IN THE BROTHERS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....46

FIGURE A1. 4. SAGE-GROUSE POPULATION TRENDS IN THE BULLY CREEK PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....47

FIGURE A1. 5. SAGE-GROUSE POPULATION TRENDS IN THE BURNS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....48

FIGURE A1. 6. SAGE-GROUSE POPULATION TRENDS IN THE COW LAKES PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....49

FIGURE A1. 7. SAGE-GROUSE POPULATION TRENDS IN THE COW VALLEY PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....50

FIGURE A1. 8. SAGE-GROUSE POPULATION TRENDS IN THE CROWLEY PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....51

FIGURE A1. 9. SAGE-GROUSE POPULATION TRENDS IN THE DIABLO PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	52
FIGURE A1. 10. SAGE-GROUSE POPULATION TRENDS IN THE DREWSEY PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	53
FIGURE A1. 11. SAGE-GROUSE POPULATION TRENDS IN THE DRY VALLEY – JACK MOUNTAIN PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	54
FIGURE A1. 12. SAGE-GROUSE POPULATION TRENDS IN THE JUNIPER MOUNTAIN PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	55
FIGURE A1. 13. SAGE-GROUSE POPULATION TRENDS IN THE LOUSE CANYON PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	56
FIGURE A1. 14. SAGE-GROUSE POPULATION TRENDS IN THE NORTH WAGONTIRE PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	57
FIGURE A1. 15. SAGE-GROUSE POPULATION TRENDS IN THE PAULINA PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	58
FIGURE A1. 16. SAGE-GROUSE POPULATION TRENDS IN THE PUEBLOS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	59
FIGURE A1. 17. SAGE-GROUSE POPULATION TRENDS IN THE SADDLE BUTTE PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	60
FIGURE A1. 18. SAGE-GROUSE POPULATION TRENDS IN THE SHEEPSHEAD PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	61
FIGURE A1. 19. SAGE-GROUSE POPULATION TRENDS IN THE SOLDIER CREEK PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	62
FIGURE A1. 20. SAGE-GROUSE POPULATION TRENDS IN THE STEENS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	63
FIGURE A1. 21. SAGE-GROUSE POPULATION TRENDS IN THE TROUT CREEKS PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	64

FIGURE A1. 22. SAGE-GROUSE POPULATION TRENDS IN THE TUCKER HILL PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	65
FIGURE A1. 23. SAGE-GROUSE POPULATION TRENDS IN THE WARNER PAC. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	66
FIGURE A1. 24. SAGE-GROUSE POPULATION TRENDS IN AREAS OUTSIDE OF OREGON’S PACS. PANEL A SHOWS THE INDEX OF SAGE-GROUSE ABUNDANCE THROUGH TIME, 1980–2024. PANEL B SHOWS THE PROPORTIONAL CHANGE IN THE INDEX OF SAGE-GROUSE ABUNDANCE FROM NADIR-TO-NADIR.....	67
<u>APPENDIX 3.</u>	
FIGURE A3. 1. GREATER SAGE-GROUSE AERIAL LEK SEARCH TRANSECTS IN SOUTHEASTERN OREGON, INCLUDING PORTIONS OF THE STEENS PAC AND ADJACENT LOW-DENSITY HABITAT, AND THE LOW-DENSITY HABITAT NORTH OF THE SHEEPSHEAD PAC, 2025.....	77
<u>APPENDIX 4.</u>	
FIGURE A4. 1. OREGON WILDLIFE MANAGEMENT UNITS WITH PERMITTED GREATER SAGE-GROUSE HUNTING AND THE DISTRIBUTION OF GREATER SAGE-GROUSE IN OREGON, 2024....	82
FIGURE A4. 2. NEST SUCCESS, PERCENT JUVENILES IN THE HARVEST, AND CHICKS PER HEN BY WMU WHERE SAGE-GROUSE HARVEST OCCURRED, OREGON, 2024. MISSING DATA IN THE BEULAH AND OWYHEE WMUS REFLECTS APPARENT NEST SUCCESS OF 0 FOR THE HENS HARVESTED IN THESE WMUS IN 2024. THIS IS AN ISSUE OF SAMPLE SIZE, AS BOTH WMUS RECORDED CHICKS IN THE HARVEST. NO WINGS WERE RECEIVED FROM THE SILVIES WMU FROM THE 2024 HUNTING SEASON.....	84
FIGURE A4. 3. OREGON SAGE-GROUSE PRODUCTION VALUES (CHICKS PER HEN; CPH) AND 32-YEAR AVERAGE (DASHED LINE; 1993–2024) ESTIMATED FROM HUNTER HARVESTED WING ANALYSES, 1993–2024.....	92
FIGURE A4. 4. ESTIMATED HATCH DATES OF MALE AND FEMALE JUVENILE SAGE-GROUSE (% HATCHED DURING PERIOD), FROM HUNTER-HARVESTED WINGS, OREGON, 2024.....	96
FIGURE A4. 5. ESTIMATED HATCH DATES OF MALE AND FEMALE JUVENILE SAGE-GROUSE (CUMULATIVE % HATCHED DURING PERIOD), FROM HUNTER-HARVESTED WINGS, OREGON, 2024.....	97

Executive Summary

During the 2025 greater sage-grouse (*Centrocercus urophasianus*) breeding season, 1,614 ground and 46 aerial lek surveys were conducted at 771 individual lek sites comprising 482 lek complexes. Surveys were conducted at 60.9% of known lek sites and 56.6% of known lek complexes in Oregon. Access to sage-grouse leks was moderate to good in most areas during the 2025 breeding season, although statewide lek survey effort was slightly below the 10-year average (2015–2024) for the proportion of known leks surveyed (10-yr. \bar{x} = 63.1%) and the proportion of known complexes surveyed (10-yr. \bar{x} = 60.8%). ODFW updated the methodology used to calculate Oregon’s sage-grouse population size estimates and trends through development of an *N*-mixture population model by the U.S. Geological Survey. Under this new model, results from the 2025 lek surveys indicate the sage-grouse spring breeding population increased by +17.8% between 2024 and 2025, to an estimated 50,912 birds (95% CI: 40,841–64,628 birds), the highest estimate recorded in Oregon since 2017. Results from 2025 lek surveys suggest Oregon’s sage-grouse populations are still trending upward in the current population cycle, and the 2025 population estimate was -2.7% below the adjusted 2003 baseline population estimate of 52,348 birds. Additionally, population abundance estimates (2017–2025) were calculated for each sage-grouse Priority Area for Conservation (PAC) using the new model. Increasing population trends between 2024 and 2025 were seen in 21 PACs, with decreasing trends in only 2 PACs, the Baker PAC (-13.3%) and Beatys PAC (-4.5%). Population increases ranged from +1.5% (Cow Lakes PAC) to +48.1% (Juniper Mountain PAC). Lastly, ODFW contracted aerial services for lek searches by helicopter in the Steens PAC and low-density habitat north of the Sheepshead PAC during spring 2025. The aerial survey documented over 19 locations where sage-grouse have not previously been known to display. Of these potential new lek locations, 7 leks were confirmed active by a ground observer in 2025 following the aerial survey, 6 in the Steens PAC and 1 lek in the low-density habitat north of the Sheepshead PAC. The remaining potential new lek locations in the Steens PAC and surrounding low-density habitat will need to be verified on the ground during the next several breeding seasons before they can be confirmed as new lek locations and added to the database.

Overview and Spring Population Monitoring Methods

Background on Lek Monitoring Efforts

Counts of male sage-grouse displaying on leks (communal breeding sites) during the spring breeding season have been used to generate indices of sage-grouse population trends since the 1940s (Patterson 1952) and remain the most widely used method to monitor sage-grouse populations across the range of the species in western North America (McCafferey et al. 2016). Monitoring of some sage-grouse leks in Oregon began in the 1940s, with survey efforts in the state beginning to increase consistently after 1980. ODFW adopted a standardized lek survey methodology in 1996, ensuring consistent data quality and allowing data comparison across the state. Lek monitoring effort increased significantly in the 2010s following the USFWS ‘warranted but precluded’ ESA listing decision for greater sage-grouse and leading up to the 2015 species review. More recently, lek survey effort increased again in 2015, following the establishment of the BLM’s Adaptive Management Strategy within the BLM 2015 Greater SageGrouse Approved Resource Management Plan (US DOI BLM OR/WA 2015). Consistently

high survey effort has been enabled by cooperative funding agreements between BLM and ODFW since 2015 which provide financial support for ODFW to hire 4–6 temporary employees to survey leks each spring, focusing on remote areas of Harney, Lake, and Malheur counties, which would otherwise receive relatively low survey effort. The agreements have also provided financial support for ODFW to reimburse volunteers within the Adopt-a-Lek (AAL) Program for their mileage and modest per diem. Lastly, the cooperative agreements have supported aerial lek searches by helicopter and aerial infrared lek counts. Based on ODFW’s aerial survey effort to date, systematic searches of core and low-density sage-grouse habitat by helicopter are the most effective way to locate new or previously unknown lekking locations and aerial infrared lek counts are the most effective way to obtain counts of males at known leks which are inaccessible on the ground.

Within previous annual reports (2013–2024), ODFW generated and reported sage-grouse breeding population estimates for each BLM District in sage-grouse range. Following adoption of ODFW’s updated Sage-Grouse Conservation Assessment and Strategy, reporting of spring population estimates transitioned from BLM Districts to Oregon’s Sage-Grouse Priority Areas for Conservation (PACs); PACs are ‘core’ sage-grouse habitats which have been geographically divided to represent biologically significant management units for Oregon’s sage-grouse populations (Figure 1). Survey effort and population trends in sage-grouse abundance will be reported at the PAC-scale in this 2025 Oregon Greater Sage-Grouse Population Monitoring Report, and in subsequent annual population monitoring reports going forward.

Lek Monitoring Methods

ODFW’s sage-grouse lek monitoring guidelines were updated in the 2025 version of the Oregon Sage-Grouse Conservation Assessment and Strategy (CAAS; Cline et al. 2025). Individual leks are those locations where two or more male sage-grouse congregate to perform their strutting display during at least two breeding seasons within a 10-year period. Lek sites must meet this requirement to be added to Oregon’s centralized sage-grouse lek database. Once a lek location has been confirmed with two or more displaying males for two years, ODFW requires only one or more displaying males present at the lek location within a 10-year period for the lek to be considered ‘occupied’. In addition to individual lek locations, ODFW identifies lek complexes as those situations where two or more individual leks are closely associated (e.g., majority of males may display at any of the known lek locations within a complex during a given breeding season or across multiple breeding seasons). This association of multiple individual leks can be identified in the field by lek observers, using data from GPS-marked males, or assumed based on geographic proximity. Typically, if two or more individual leks are within 1.6 km (1 mi) of the next closest known lek, these leks are grouped and considered a single lek complex. Importantly, all individual leks within a lek complex must be surveyed on the same morning to avoid biases associated with the intra-season movements of males within a complex.

Sage-grouse leks and lek complexes in southeastern Oregon are monitored between 15 March and 30 April, annually, to obtain counts of breeding male sage-grouse. In a collaborative effort, biologists with ODFW, BLM, USFWS, Burns Paiute Tribe, and Oregon State University, as well as volunteers enrolled in the ODFW Adopt-a-Lek Program (Appendix II), visit leks each spring to count all male and female sage-grouse visible on a lek. Due to variability in male lek attendance throughout the breeding season, a subset of lek complexes is counted up to 3 times per season, with individual complex counts separated by at least 7 days.

Lek counts are used to assess sage-grouse population size and trends; the following protocol must be followed to ensure unbiased estimates:

- Male sage-grouse begin attending leks in early March and often continue through early May. Lek counts should be conducted only between 15 March–30 April each year to coincide with peak female attendance.
- Lek counts must be conducted only from 30 minutes before sunrise to 2 hours after sunrise. Lek counts conducted outside of this window are not included in the data analyses to estimate Oregon’s annual sage-grouse abundance (Appendix 2.). Detection probability of males declines significantly beyond 45 minutes post-sunrise, so whenever possible, lek surveys should be completed during this timeframe.
- Lek counts should be conducted during mornings with clear to partly cloudy skies, little to no wind, and no precipitation. Lek counts should not be conducted if weather conditions (i.e., wind gusts >15 mph (24 km/h), rainy or snowy conditions) or disturbance (i.e., humans, livestock, predators, other wildlife, etc.) precludes normal display activities.
- For unbiased estimates of population size and trends, a minimum of 50% of the ‘occupied’ or ‘occupied-pending’ lek complexes (see below for definitions) within a geographic scale of interest (i.e., PAC, WMU, BLM District, statewide, etc.) should be counted at least 2 times, annually. Lek complexes should be counted at 7–14-day intervals. If a geographic scale of interest contains <10 active or active pending leks, a minimum of 80% of these leks should be counted at least once annually.
- All known ‘occupied’ or ‘occupied-pending’ lek complexes on accessible public or private lands should be surveyed a minimum of every 3 years.
- All known ‘unoccupied’ or ‘unoccupied-pending’ lek complexes on accessible public or private lands should be surveyed a minimum of every 6 years, ideally during the peak of a population cycle not during the trough of a population cycle.

Ground Surveys

Sage-grouse lek counts should primarily be conducted from the ground when the lek location is accessible. Ground surveys should be conducted by a trained observer viewing the lek through a spotting scope or binoculars from a concealed location, generally >200 m (>656 ft) from the lek to avoid disturbing the sage-grouse. The observer should record survey information (i.e., lek identifier and name, date, time, weather conditions) at the beginning of the survey; data may be recorded electronically or on paper datasheets in the field. The observer should count the sage-grouse on the lek from right to left, or left to right, recording the number of males, females, and sage-grouse of unknown sex, and report each separately. The observer should survey each lek

until the maximum male count recorded has remained the same for two or more consecutive counts, and for a minimum of 15 minutes.

Presence-Absence Checks

Presence-absence sage-grouse lek checks report lek activity when a lek count cannot be obtained, documenting the lek status as active/detected or not detected during a given breeding season. For example, if an observer visits a known lek location outside of the dawn lek count timeframe or when weather conditions preclude a lek count and the observer detects male sage-grouse at the lek, this lek may be active during the given breeding season and an observer should return on a subsequent morning to conduct a lek count. Additional signs of activity include documentation of fresh sage-grouse droppings, feathers, or tracks. Importantly, if sage-grouse or sage-grouse sign are not observed at a known lek when the lek is visited outside of the lek count timeframe or viewed under conditions that preclude a lek count, the lek should not be reported as ‘inactive’ during the given breeding season. Rather, the observer should report neither sage-grouse nor sage-grouse sign were detected during the presence-absence lek check. Although presence-absence checks for activity at sage-grouse leks may provide some useful information (e.g., prioritization for subsequent lek count), the structure of Oregon’s centralized sage-grouse lek database precludes the utility of presence-absence checks. Oregon tracks annual lek activity status based on lek counts and does not incorporate presence-absence checks into annual reporting of population status or trend at any geographic scale. Presence-absence lek checks should be collected opportunistically and reported but should not be prioritized over lek counts.

Table 1. Definitions of ODFW’s Sage-Grouse Lek Conservation Statuses, 2025.

Lek Status	10-year survey period	11–20-year period	21–30-year period
Occupied	1+ males counted during 2+ surveys		
Occupied-Pending	1+ males counted during 1 survey		
	No surveys conducted	1+ males counted during the most recent survey	
Unoccupied-Pending	Only 1 survey with 0 males observed	And 1+ males counted during 1+ surveys	
	No surveys conducted	And 1+ males counted during 1+ surveys	
	No surveys conducted	And no surveys conducted	And 1+ males counted during 1+ surveys
Unoccupied	0 males counted during 2+ surveys	And 1+ males counted during 1+ surveys	
Historic	0 males counted during 1+ surveys		
	0 males recorded within the last 30 years		

Lek Data Analyses

Previous versions of the annual Oregon Greater Sage-Grouse Population Monitoring Report (2011–2024) presented Oregon’s sage-grouse population size estimates and population trends obtained from the breeding population analysis described in Hagen (2011). This methodology was updated in 2025 when ODFW contracted with USGS to develop a new sage-grouse population model (*N*-Mixture model) for Oregon; see ODFW CAAS (Cline et al. 2025) Appendix 2: Population Analysis Methodology for the full report.

The updated model accounts for imperfect detection of male sage-grouse attending leks and provides unbiased estimates of sage-grouse abundance at large (Oregon range-wide) and smaller (PAC) spatial scales. Although the sage-grouse population trend estimates from the two models are similar, point estimates of population size are different. Three key differences between the previous and new population models explain these discrepancies:

- 1) Oregon’s previous population model did not account for the specific variables influencing detection of sage-grouse during a given survey (i.e., time since sunrise; Figure 2A), within a survey season (i.e., Julian date; Figure 2B), or the interannual variation in detection probability among years, each of which the new model estimates and incorporates. Without accounting for variability in detection probability of males attending known leks, the previous model functioned as an index of minimum population size, where the new model estimates true abundance of sage-grouse.
- 2) The new population model incorporates variability in male attendance to leks within a breeding season to account for the ‘non-lek mating’ cohort, which the previous population model did not consider. Within the new model, the age-weighted average male lek attendance rate was defined as 0.58 (95% CI = 0.46–0.71; Wann et al. 2019).
- 3) The new model adjusts for presently unknown leks on the landscape by considering those previously unknown leks added to the database annually, 1941–2025. There have been over 100 new leks discovered in Oregon and added to the statewide lek database over the past 10 years. Under ODFW’s previous sage-grouse population model, new leks were mostly unaccounted for, resulting in large apparent population increases following their addition to the lek database.

The *N*-Mixture model requires repeated counts (e.g., multiple surveys per season) to provide unbiased estimates of sage-grouse abundance, and the years from 2017–present have a sufficient sample size of repeat surveys to run the model (Figure 5). To estimate the statewide sage-grouse population abundance during the period from 1980–2016, we fit a general linear model using the 2017–present population estimates from the new model and the previous model [New Model Estimates \sim 1 + Previous Model Estimates; $p = 0.013$, $r\text{-sq.} = 0.61$]. We used the results of this general linear model to predict the updated statewide population abundance during 1980–2016 from the population estimates of the previous model during this timeframe (Figure 6).




Figure 2. The effect of time since sunrise (Panel A) and date (Panel B) on the probability of detecting male sage-grouse attending leks in eastern Oregon, 2017–2025.

Lek Monitoring Effort and Population Estimates

Statewide

ODFW and partners completed 1,614 ground surveys and 46 aerial counts during mid-March through April 2025. Surveys were conducted at 771 leks comprising 482 lek complexes. Of the 1,267 individual leks, and 851 lek complexes known to exist or have existed in the state, 60.9% and 56.6%, respectively, were surveyed during 2025 (Figure 3). On average, each lek was surveyed 2.15 times. Statewide lek survey effort during the 2025 breeding season was slightly below the 10-year average (2015–2024) for the proportion of known leks surveyed (10-yr. \bar{x} = 63.1%) and the proportion of known complexes surveyed (10-yr. \bar{x} = 60.8%). Access to sage-grouse leks was moderate to good in most areas, so lower survey effort this spring was driven by other factors in addition to access and road conditions.




Figure 3. Number of sage-grouse leks and lek complexes known to exist in Oregon since 1980, and the number of leks and lek complexes surveyed each year, 1980–2025.

Dedicated aerial surveys (Appendix III) and incidental observations during ground surveys help expand knowledge of sage-grouse distribution in Oregon. ODFW contracted aerial services for lek searches by helicopter in the Steens PAC and low-density habitat north of the Sheepshead PAC during spring 2025. The aerial survey documented over 19 locations where sage-grouse have not previously been known to display. Of these potential new lek locations, 7 leks were confirmed active by a ground observer in 2025 following the aerial survey, 6 in the Steens PAC and 1 lek in the low-density habitat north of the Sheepshead PAC. The remaining potential new lek locations in the Steens PAC and surrounding low-density habitat will need to be verified on

the ground during the next several breeding seasons before they can be confirmed as new lek locations and added to the database. Four previously unknown leks discovered during 2024 aerial surveys were checked and confirmed during the 2025 breeding season, 3 new leks in the Steens PAC and 1 new lek in the Pueblos PAC. Additionally, two previously unknown leks discovered during the 2022 aerial survey were confirmed on the ground during the 2025 lek survey season and added to the statewide lek database, one lek in the Louse Canyon PAC and one lek in the Trout Creeks PAC. There are several additional potential new lek locations in these PACs which still need to be confirmed on the ground before they can be added to the statewide lek database. Finally, two previously unknown leks discovered by Vale District BLM staff in the Bully Creek PAC were confirmed active and added to the database in 2025.




Figure 4. Number of sage-grouse lek surveys conducted per year in Oregon (blue line), and the average number of surveys conducted at each surveyed lek by year (orange line), 1980–2025.

The estimated spring greater sage-grouse population in Oregon during 2025 was 50,912 birds (95% CI: 40,841–64,628 birds), a 17.8% increase from 2024 (2024 estimate = 43,233 birds [95% CI: 34,914–54,754 individuals]; Figure 5). The 2025 population estimate was -2.7% below the 2003 baseline population estimate of 52,348 individuals. The statewide population estimate for 2025 is the highest recorded in Oregon since 2017, which was the most recent peak in Oregon’s sage-grouse population cycle, and suggests Oregon’s sage-grouse populations are still trending upward in the current population cycle. The most recent population nadir (trough) was evident in 2019 when the statewide population was estimated at 32,553 sage-grouse, the lowest statewide estimate during the 46-year period from 1980–2025 (Figure 6). Population trends (i.e., changes in population size) should be interpreted from complete population cycles using the population

nadirs (Coates et al. 2021). During the 1980–2025 timeframe, five nadirs exist in Oregon’s statewide sage-grouse population cycles, in years 1985, 1996, 2008, 2013, and 2019. Comparing trends for each of these four previous population nadirs to the most recent nadir, Oregon’s sage-grouse populations, statewide, have declined by -32.9% (1985–2019), -20.0% (1996–2019), -11.9% (2008–2019), and -13.8% (2013–2019).




Figure 5. *N*-mixture model estimates of sage-grouse abundance within the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.




Figure 6. Oregon statewide sage-grouse population estimates, 1980–2025. The solid black line represents the abundance estimates from the new *N*-mixture model, and the dashed gray line represents the population estimates from ODFW’s previous model. The population goal is set at the 2003 level, representing the baseline for Oregon’s statewide sage-grouse population size.

Abundance Results (2025), PAC-Scale

Habitat management for sage-grouse in Oregon often occurs at the scale of individual Priority Areas for Conservation (PAC), or at even smaller scales within a PAC, so knowing the abundance of sage-grouse within each of Oregon’s sage-grouse PACs is important for the management and conservation of the species. Sage-grouse population abundance estimates (2017–2025) from ODFW’s new *N*-Mixture model are presented in this section for each PAC. Of Oregon’s 23 PACs, 8 PACs make up the majority (>70%) of Oregon’s sage-grouse population by abundance: Beatys PAC (14.0%), Steens PAC (10.5%), Louse Canyon PAC (10.4%), Warner PAC (8.3%), Trout Creeks PAC (8.0%), Bully Creek PAC (6.8%), Crowley PAC (6.7%), and Pueblos PAC (6.6%; Table 2). Increasing population trends between 2024 and 2025 were seen in 21 PACs, with decreasing trends in only 2 PACs, the Baker PAC (-13.3%) and Beatys PAC (-4.5%). Population increases ranged from +1.5% (Cow Lakes PAC) to +48.1% (Juniper Mountain PAC).

Note, estimates of true abundance from the *N*-Mixture model are unavailable at the PAC-scale prior to the year 2017 due to an inadequate sample size of repeat surveys conducted within

survey seasons before 2017. However, estimates of relative sage-grouse abundance are available through the USGS Rangewide Population Monitoring Framework (Ver. 4, Coates et al. 2025), detailed in Appendix I.

Table 2. Estimated sage-grouse abundance (2025) and percent of the 2025 spring breeding sage-grouse population within each of Oregon’s Priority Areas for Conservation (PACs).

Priority Area for Conservation (PAC)	Abundance, 2025	Abundance SD ¹ , 2025	Trend ² (%), 2025	% of 2025 Population
Baker	475	108.6	-13.3	0.9
Beatys	7,155	1110.7	-4.5	14.0
Brothers	1,343	177.1	31.0	2.6
Bully Creek	3,463	521.6	11.5	6.8
Burns	178	36.3	12.8	0.3
Cow Lakes	935	159.0	1.5	1.8
Cow Valley	1,808	324.2	4.4	3.5
Crowley	3,427	450.1	36.7	6.7
Diablo	91	59.1	19.4	0.2
Drewsey	1,865	266.3	21.9	3.7
Dry Valley/Jack Mountain	856	165.1	16.3	1.7
Juniper Mountain	386	115.6	48.1	0.8
Louse Canyon	5,288	873.8	28.0	10.4
North Wagontire	464	67.6	14.2	0.9
Paulina	2,148	331.6	24.6	4.2
Pueblos	3,379	441.2	23.5	6.6
Saddle Butte	674	199.4	11.0	1.3
Sheepshead	196	86.2	18.1	0.4
Soldier Creek	2,071	403.6	12.5	4.1
Steens	5,368	728.4	33.3	10.5
Trout Creeks	4,101	573.5	23.8	8.0
Tucker Hill	341	49.2	38.2	0.7
Warner	4,218	548.6	27.1	8.3

¹Standard deviation of the abundance estimate.

²Percent change (trend) in estimated abundance between 2024 and 2025.

Baker PAC

The Baker PAC is an area of approximately 115,262 ha (284,815 ac), located mostly within Baker County; a small portion of the PAC falls within Union County. Sixty-two leks are known to exist in the Baker PAC. As of 2025, the conservation statuses of these leks are: 16 occupied, 4 occupied-pending, 3 unoccupied-pending, 18 unoccupied, and 21 historic leks.

During 2025, 63 lek surveys were conducted at 26 leks within the Baker PAC; 20 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Baker PAC was counted 2.42 times. Estimated sage-grouse abundance in the Baker PAC in 2025 was 475 birds (95% CI: 324–740 birds), down -13.3% from the 548 birds estimated in the PAC in 2024 (Figure 7). The Baker PAC population makes up 0.9% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Baker PAC was -22.8% during 2014–2023, the mid-term population trend (2 oscillations) was -79.6% during 2001–2023, and the long-term population trend (4 oscillations) was -81.5% during 1985–2023 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Baker PAC are available in Appendix I (Figure A1.1).




Figure 7. *N*-mixture model estimates of sage-grouse abundance within Baker Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Beatys PAC

The Beatys PAC is an area of approximately 337,541 ha (834,073 ac), located within Lake County and Harney County, and contains 166 known leks. As of 2025, the conservation statuses of these leks are: 85 occupied, 21 occupied-pending, 17 unoccupied, and 43 historic leks.

During 2025, 267 lek surveys were conducted at 103 leks within the Beatys PAC; 84 (79%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Beatys PAC was counted 2.59 times. Estimated sage-grouse abundance in the Beatys PAC in 2025 was 7,154 birds (95% CI: 5,345–9,637 birds), down -4.5% from the 7,488 birds estimated in the PAC in 2024 (Figure 8). The Beatys PAC population makes up 14.0% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Beatys PAC was -17.4% during 2013–2019, the mid-term population trend (2 oscillations) was -72.7% during 2002–2019, and the long-term population trend (4 oscillations) was -76.5% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Beatys PAC are available in Appendix I (Figure A1.2).




Figure 8. *N*-mixture model estimates of sage-grouse abundance within Beatys Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Brothers PAC

The Brothers PAC is an area of approximately 156,589 ha (386,936 ac), located mostly within Deschutes County; a portion of the PAC falls within Crook County and Lake County. Fifty leks are known to exist in the Brothers PAC. As of 2025, the conservation statuses of these leks are: 30 occupied, 3 occupied-pending, 8 unoccupied, and 9 historic leks.

During 2025, 131 lek surveys were conducted at 43 leks within the Brothers PAC; 33 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Brothers PAC was counted 3.05 times. Estimated sage-grouse abundance in the Brothers PAC in 2025 was 1,343 birds (95% CI: 1,059–1,747 birds), up +31.0% from the 1,025 birds estimated in the PAC in 2024 (Figure 9). The Brothers PAC population makes up 2.6% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Brothers PAC was -58.7% during 2009–2019, the mid-term population trend (2 oscillations) was -53.5% during 2003–2019, and the long-term population trend (4 oscillations) was -80.9% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Brothers PAC are available in Appendix I (Figure A1.3).




Figure 9. *N*-mixture model estimates of sage-grouse abundance within Brothers Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Bully Creek PAC

The Bully Creek PAC is an area of approximately 201,538 ha (498,006 ac), located mostly within Malheur County; small portions of the PAC fall within Grant County and Harney County. Fifty-nine leks are known to exist in the Bully Creek PAC. As of 2025, the conservation statuses of these leks are: 38 occupied, 4 occupied-pending, 7 unoccupied, and 10 historic leks.

During 2025, 88 lek surveys were conducted at 44 leks within the Bully Creek PAC; 38 (90.5%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Bully Creek PAC was counted 2.0 times. Estimated sage-grouse abundance in the Bully Creek PAC in 2025 was 3,463 birds (95% CI: 2,628–4,660 birds), up +11.5% from the 3,106 birds estimated in the PAC in 2024 (Figure 10). The Bully Creek PAC population makes up 6.8% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Bully Creek PAC was -31.1% during 2014–2020, the mid-term population trend (3 oscillations) was -35.8% during 2001–2020, and the long-term population trend (5 oscillations) was -20.9% during 1985–2020 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Bully Creek PAC are available in Appendix I (Figure A1.4).




Figure 10. *N*-mixture model estimates of sage-grouse abundance within Bully Creek Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Burns PAC

The Burns PAC is an area of approximately 31,098 ha (76,845 ac), located within Harney County, and contains 5 known leks. As of 2025, the conservation statuses of these leks are: 3 occupied and 2 historic leks.

During 2025, 8 lek surveys were conducted at the 3 occupied leks within the Burns PAC. On average, each surveyed lek in the Burns PAC was counted 2.67 times. Estimated sage-grouse abundance in the Burns PAC in 2025 was 178 birds (95% CI: 129–268 birds), up +12.8% from the 158 birds estimated in the PAC in 2024 (Figure 11). The Burns PAC population makes up 0.3% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Burns PAC was -32.7% during 2014–2020, the mid-term population trend (3 oscillations) was -49.8% during 2002–2020, and the long-term population trend (5 oscillations) was -66.9% during 1985–2020 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Burns PAC are available in Appendix I (Figure A1.5).




Figure 11. *N*-mixture model estimates of sage-grouse abundance within Burns Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Cow Lakes PAC

The Cow Lakes PAC is an area of approximately 141,890 ha (350,613 ac), located within Malheur County, and contains 63 known leks. As of 2025, the conservation statuses of these leks are: 26 occupied, 6 occupied-pending, 16 unoccupied, and 15 historic leks.

During 2025, 91 lek surveys were conducted at 47 leks within the Cow Lakes PAC; 31 (96.9%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Cow Lakes PAC was counted 1.94 times. Estimated sage-grouse abundance in the Cow Lakes PAC in 2025 was 935 birds (95% CI: 698–1,306 birds), up +1.5% from the 922 birds estimated in the PAC in 2024 (Figure 12). The Cow Lakes PAC population makes up 1.8% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Cow Lakes PAC was -29.6% during 2013–2022, the mid-term population trend (3 oscillations) was -68.7% during 2001–2022, and the long-term population trend (5 oscillations) was -85.4% during 1985–2022 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Cow Lakes PAC are available in Appendix I (Figure A1.6).




Figure 12. *N*-mixture model estimates of sage-grouse abundance within Cow Lakes Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Cow Valley PAC

The Cow Valley PAC is an area of approximately 164,097 ha (405,489 ac), located within northern Malheur County and southern Baker County, and contains 59 known leks. As of 2025, the conservation statuses of these leks are: 36 occupied, 7 occupied-pending, 2 unoccupied, and 14 historic leks.

During 2025, 68 lek surveys were conducted at 33 leks within the Cow Lakes PAC; 28 (65.1%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Cow Valley PAC was counted 2.06 times. Estimated sage-grouse abundance in the Cow Valley PAC in 2025 was 1,808 birds (95% CI: 1,297–2,559 birds), up +4.4% from the 1,732 birds estimated in the PAC in 2024 (Figure 13). The Cow Valley PAC population makes up 3.5% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Cow Valley PAC was +1.9% during 2014–2019, the mid-term population trend (3 oscillations) was -21.8% during 2001–2019, and the long-term population trend (5 oscillations) was -3.3% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Cow Valley PAC are available in Appendix I (Figure A1.7).




Figure 13. *N*-mixture model estimates of sage-grouse abundance within Cow Valley Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Crowley PAC

The Crowley PAC is an area of approximately 238,168 ha (588,519 ac), located mostly within Malheur County; a small portion of the PAC falls within Harney County. Fifty-three leks are known to exist in the Crowley PAC. As of 2025, the conservation statuses of these leks are: 33 occupied, 2 occupied-pending, 6 unoccupied, and 12 historic leks.

During 2025, 110 lek surveys were conducted at 49 leks within the Crowley PAC; 35 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Crowley PAC was counted 2.24 times. Estimated sage-grouse abundance in the Crowley PAC in 2025 was 3,427 birds (95% CI: 2,687–4,410 birds), up +36.7% from the 2,507 birds estimated in the PAC in 2024 (Figure 14). The Crowley PAC population makes up 6.7% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Crowley PAC was -16.9% during 2013–2021, the mid-term population trend (3 oscillations) was -38.1% during 2002–2021, and the long-term population trend (5 oscillations) was -55.5% during 1984–2021 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Crowley PAC are available in Appendix I (Figure A1.8).




Figure 14. *N*-mixture model estimates of sage-grouse abundance within Crowley Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Diablo PAC

The Diablo PAC is an area of approximately 22,852 (56,469 ac), located within Lake County, and contains 3 known leks. As of 2025, the conservation statuses of these leks are: 1 occupied and 2 historic leks.

During 2025, 2 lek surveys were conducted at the 1 occupied lek within the Diablo PAC. Estimated sage-grouse abundance in the Diablo PAC in 2025 was 91 birds (95% CI: 49–257 birds), up +19.3% from the 76 birds estimated in the PAC in 2024 (Figure 15). The Diablo PAC population makes up 0.2% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Diablo PAC was -42.6% during 2013–2021, the mid-term population trend (3 oscillations) was -60.2% during 2002–2021, and the long-term population trend (5 oscillations) was -71.8% during 1985–2021 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Diablo PAC are available in Appendix I (Figure A1.9).




Figure 15. *N*-mixture model estimates of sage-grouse abundance within Diablo Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Drewsey PAC

The Drewsey PAC is an area of approximately 178,879 ha (442,015 ac), located mostly within Harney County; a small portion of the PAC falls within Malheur County. Fifty-one leks are known to exist in the Drewsey PAC. As of 2025, the conservation statuses of these leks are: 39 occupied, 4 occupied-pending, 1 unoccupied, and 7 historic leks.

During 2025, 79 lek surveys were conducted at 38 leks within the Drewsey PAC; 36 (83.7%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Drewsey PAC was counted 2.08 times. Estimated sage-grouse abundance in the Drewsey PAC in 2025 was 1,865 birds (95% CI: 1,441–2,476 birds), up +21.9% from the 1,530 birds estimated in the PAC in 2024 (Figure 16). The Drewsey PAC population makes up 3.7% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Drewsey PAC was -31.3% during 2013–2019, the mid-term population trend (3 oscillations) was -43.7% during 2002–2019, and the long-term population trend (5 oscillations) was -73.3% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Drewsey PAC are available in Appendix I (Figure A1.10).




Figure 16. *N*-mixture model estimates of sage-grouse abundance within Drewsey Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Dry Valley – Jack Mountain PAC

The Dry Valley – Jack Mountain PAC is an area of approximately 80,749 ha (199,532 ac), located within Harney County, and contains 12 known leks. As of 2025, the conservation statuses of these leks are: 5 occupied, 1 occupied-pending, 1 unoccupied, and 5 historic leks.

During 2025, 13 lek surveys were conducted at 7 leks within the Dry Valley – Jack Mountain PAC; 6 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Dry Valley – Jack Mountain PAC was counted 1.86 times. Estimated sage-grouse abundance in the Dry Valley – Jack Mountain PAC in 2025 was 856 birds (95% CI: 641–1,275 birds), up +16.3% from the 736 birds estimated in the PAC in 2024 (Figure 17). The Dry Valley – Jack Mountain PAC population makes up 1.7% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Dry Valley – Jack Mountain PAC was stable (0.0%) during 2015–2019, the mid-term population trend (3 oscillations) was -71.2% during 2002–2019, and the long-term population trend (5 oscillations) was -85.7% during 1984–2019 (Table A1.1) Figures illustrating the long-term trends (1980–2024) for the Dry Valley/Jack Mountain PAC are available in Appendix I (Figure A1.11).




Figure 17. *N*-mixture model estimates of sage-grouse abundance within Dry Valley – Jack Mountain Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Juniper Mountain PAC

The Juniper Mountain PAC is an area of approximately 55,493 ha (137,125 ac), located within Harney County and Lake County, and contains 11 known leks. As of 2025, the conservation statuses of these leks are: 4 occupied, 1 occupied-pending, and 6 historic leks.

During 2025, 3 lek surveys were conducted at 1 lek within the Juniper Mountain PAC; 1 (20%) of the occupied or occupied-pending leks was surveyed one or more times in 2025. On average, each surveyed lek in the Juniper Mountain PAC was counted 3.0 times. Estimated sage-grouse abundance in the Juniper Mountain PAC in 2025 was 386 birds (95% CI: 249–696 birds), up +48.0% from the 261 birds estimated in the PAC in 2024 (Figure 18). The Juniper Mountain PAC population makes up 0.8% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Juniper Mountain PAC was +10.1% during 2013–2019, the mid-term population trend (3 oscillations) was +16.6% during 1996–2019, and the long-term population trend (4 oscillations) was -32.4% during 1984–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Juniper Mountain PAC are available in Appendix I (Figure A1.12).




Figure 18. *N*-mixture model estimates of sage-grouse abundance within Juniper Mountain Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Louse Canyon PAC

The Louse Canyon PAC is an area of approximately 222,584 ha (550,012 ac), located within Malheur County, and contains 62 known leks. As of 2025, the conservation statuses of these leks are: 32 occupied, 6 occupied-pending, 7 unoccupied, and 17 historic leks.

During 2025, 58 lek surveys were conducted at 42 leks within the Louse Canyon PAC; 34 (89.5%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Louse Canyon PAC was counted 1.38 times. Estimated sage-grouse abundance in the Louse Canyon PAC in 2025 was 5,287 birds (95% CI: 3,938–7,366 birds), up +28.0% from the 4,132 birds estimated in the PAC in 2024 (Figure 19). The Louse Canyon PAC population makes up 10.4% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Louse Canyon PAC was -21.2% during 2014–2019, the mid-term population trend (2 oscillations) was -25.0% during 1996–2019, and the long-term population trend (3 oscillations) was -56.8% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Louse Canyon PAC are available in Appendix I (Figure A1.13).




Figure 19. *N*-mixture model estimates of sage-grouse abundance within Louse Canyon Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

North Wagontire PAC

The North Wagontire PAC is an area of approximately 110,676 ha (273,484 ac), located mostly within Lake County; a portion of the PAC falls within Deschutes County and Harney County. Nineteen leks are known to exist in the North Wagontire PAC. As of 2025, the conservation statuses of these leks are: 10 occupied, 1 occupied-pending, 1 unoccupied, and 7 historic leks.

During 2025, 32 lek surveys were conducted at 12 leks within the North Wagontire PAC; 11 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the North Wagontire PAC was counted 2.67 times. Estimated sage-grouse abundance in the North Wagontire PAC in 2025 was 463 birds (95% CI: 361–623 birds), up +14.1% from the 406 birds estimated in the PAC in 2024 (Figure 20). The North Wagontire PAC population makes up 0.9% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the North Wagontire PAC was -33.7% during 2013–2023, the mid-term population trend (2 oscillations) was -49.5% during 2002–2023, and the long-term population trend (4 oscillations) was -73.7% during 1985–2023 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the North Wagontire PAC are available in Appendix I (Figure A1.14).




Figure 20. *N*-mixture model estimates of sage-grouse abundance within North Wagontire Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Paulina PAC

The Paulina PAC is an area of approximately 225,290 ha (556,698 ac), located mostly within Crook County; a portion of the PAC falls within Deschutes County, Harney County, and Lake County. Sixty-eight leks are known to exist in the Paulina PAC. As of 2025, the conservation statuses of these leks are: 35 occupied, 8 occupied-pending, 5 unoccupied-pending, 8 unoccupied, and 12 historic leks.

During 2025, 55 lek surveys were conducted at 38 leks within the Paulina PAC; 30 (69.8%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Paulina PAC was counted 1.45 times. Estimated sage-grouse abundance in the Paulina PAC in 2025 was 2,148 birds (95% CI: 1,610–2,874 birds), up +24.6% from the 1,725 birds estimated in the PAC in 2024 (Figure 21). The Paulina PAC population makes up 4.2% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Paulina PAC was -15.4% during 2013–2023, the mid-term population trend (3 oscillations) was -51.7% during 2002–2023, and the long-term population trend (5 oscillations) was -60.3% during 1985–2023 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Paulina PAC are available in Appendix I (Figure A1.15).




Figure 21. *N*-mixture model estimates of sage-grouse abundance within Paulina Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Pueblos PAC

The Pueblos PAC is an area of approximately 143,527 ha (354,659 ac), located within Harney County, and contains 47 known leks. As of 2025, the conservation statuses of these leks are: 29 occupied, 2 occupied-pending, 10 unoccupied, and 6 historic leks.

During 2025, 44 lek surveys were conducted at 23 leks within the Pueblos PAC; 22 (68.8%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Pueblos PAC was counted 1.91 times. Estimated sage-grouse abundance in the Pueblos PAC in 2025 was 3,379 birds (95% CI: 2,655–4,402 birds), up +23.4% from the 2,737 birds estimated in the PAC in 2024 (Figure 22). The Pueblos PAC population makes up 6.6% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend in the Pueblos PAC was -5.8% during 2019–2022 (1 oscillation) and -31.8% during 2013–2022 (2 oscillations). The mid-term population trend (3 oscillations) was -31.4% during 2009–2022, and the long-term population trend (5 oscillations) was -53.2% during 1985–2022 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Pueblos PAC are available in Appendix I (Figure A1.16).




Figure 22. *N*-mixture model estimates of sage-grouse abundance within Pueblos Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Saddle Butte PAC

The Saddle Butte PAC is an area of approximately 43,189 ha (106,722 ac), located in Malheur County, and contains 5 known leks. As of 2025, the conservation statuses of these leks are: 4 occupied, and 1 occupied-pending leks.

During 2025, 17 lek surveys were conducted at 5 leks within the Saddle Butte PAC; 5 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Saddle Butte PAC was counted 3.4 times. Estimated sage-grouse abundance in the Saddle Butte PAC in 2025 was 674 birds (95% CI: 382–1,182 birds), up +11.0% from the 607 birds estimated in the PAC in 2024 (Figure 23). The Saddle Butte PAC population makes up 1.3% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Saddle Butte PAC was +55.5% during 2013–2020, the mid-term population trend (3 oscillations) was +40.5% during 2002–2020, and the long-term population trend (5 oscillations) was stable (+0.0%) during 1985–2020 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Saddle Butte PAC are available in Appendix I (Figure A1.17).




Figure 23. *N*-mixture model estimates of sage-grouse abundance within Saddle Butte Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Sheepshead PAC

The Sheepshead PAC is an area of approximately 20,084 ha (49,627 ac), located within Harney County and Malheur County, and contains 4 known leks. As of 2025, the conservation statuses of these leks are: 2 occupied, 1 unoccupied, and 1 historic lek.

During 2025, 4 ground surveys and 1 aerial lek survey were conducted at 3 leks within the Sheepshead PAC; 2 (100%) of the occupied leks were surveyed one or more times in 2025. On average, each surveyed lek in the Sheepshead PAC was counted 1.67 times. Estimated sage-grouse abundance in the Sheepshead PAC in 2025 was 196 birds (95% CI: 117–437 birds), up +18.1% from the 166 birds estimated in the PAC in 2024 (Figure 24). The Sheepshead PAC population makes up 0.4% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Sheepshead PAC was -24.2% during 2013–2021, the mid-term population trend (3 oscillations) was -48.2% during 2002–2021, and the long-term population trend (5 oscillations) was -65.4% during 1985–2021 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Sheepshead PAC are available in Appendix I (Figure A1.18)




Figure 24. *N*-mixture model estimates of sage-grouse abundance within Sheepshead Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Soldier Creek PAC

The Soldier Creek PAC is an area of approximately 163,915 ha (405,039 ac), located in Malheur County, and contains 67 known leks. As of 2025, the conservation statuses of these leks are: 35 occupied, 5 occupied-pending, 12 unoccupied, and 15 historic leks.

During 2025, 96 lek surveys were conducted at 45 leks within the Soldier Creek PAC; 34 (85.0%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Soldier Creek PAC was counted 2.13 times. Estimated sage-grouse abundance in the Soldier Creek PAC in 2025 was 2,071 birds (95% CI: 1,448–3,028 birds), up +12.5% from the 1,840 birds estimated in the PAC in 2024 (Figure 25). The Soldier Creek PAC population makes up 4.1% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Soldier Creek PAC was -37.9% during 2013–2020, the mid-term population trend (3 oscillations) was -52.2% during 1996–2020, and the long-term population trend (4 oscillations) was -75.2% during 1985–2020 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Soldier Creek PAC are available in Appendix I (Figure A1.19).




Figure 25. *N*-mixture model estimates of sage-grouse abundance within Soldier Creek Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Steens PAC

The Steens PAC is an area of approximately 222,216 ha (549,101 ac), located mostly within Harney County; a small portion of the PAC falls within Malheur County. Fifty-one leks are known to exist in the Steens PAC. As of 2025, the conservation statuses of these leks are: 36 occupied, 5 occupied-pending, 1 unoccupied, and 9 historic leks.

During 2025, 61 ground and 41 aerial lek surveys were conducted at 48 leks within the Steens PAC; 41 (100%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Steens PAC was counted 2.10 times. Estimated sage-grouse abundance in the Steens PAC in 2025 was 5,368 birds (95% CI: 4,189–6,992 birds), up +33.3% from the 4,027 birds estimated in the PAC in 2024 (Figure 26). The Steens PAC population makes up 10.5% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Steens PAC was -10.2% during 2013–2020, the mid-term population trend (3 oscillations) was -10.4% during 1996–2020, and the long-term population trend (4 oscillations) was -45.4% during 1985–2020 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Steens PAC are available in Appendix I (Figure A1.20).




Figure 26. *N*-mixture model estimates of sage-grouse abundance within Steens Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Trout Creeks PAC

The Trout Creeks PAC is an area of approximately 193,032 ha (476,988 ac), located within Harney County and Malheur County, and contains 104 known leks. As of 2025, the conservation statuses of these leks are: 53 occupied, 9 occupied-pending, 21 unoccupied, and 21 historic leks.

During 2025, 155 lek surveys were conducted at 76 leks within the Trout Creeks PAC; 53 (85.5%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Trout Creeks PAC was counted 2.04 times. Estimated sage-grouse abundance in the Trout Creeks PAC in 2025 was 4,101 birds (95% CI: 3,161–5,362 birds), up +23.8% from the 3,312 birds estimated in the PAC in 2024 (Figure 27). The Trout Creeks PAC population makes up 8.0% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Trout Creeks PAC was +7.1% during 2013–2021, the mid-term population trend (3 oscillations) was -18.4% during 2002–2021, and the long-term population trend (5 oscillations) was -49.3% during 1985–2021 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Trout Creeks PAC are available in Appendix I (Figure A1.21).




Figure 27. *N*-mixture model estimates of sage-grouse abundance within Trout Creeks Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Tucker Hill PAC

The Tucker Hill PAC is an area of approximately 12,482 ha (30,844 ac), located within Lake County, and contains 6 known leks. As of 2025, the conservation statuses of these leks are: 4 occupied and 2 historic leks.

During 2025, 12 lek surveys were conducted at the 4 occupied leks within the Tucker Hill PAC. On average, each surveyed lek in the Tucker Hill PAC was counted 3.0 times. Estimated sage-grouse abundance in the Tucker Hill PAC in 2025 was 341 birds (95% CI: 266–455 birds), up +38.2% from the 247 birds estimated in the PAC in 2024 (Figure 28). The Tucker Hill PAC population makes up 0.7% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Tucker Hill PAC was -8.4% during 2013–2023, the mid-term population trend (3 oscillations) was +3.6% during 2001–2023, and the long-term population trend (5 oscillations) was +61.1% during 1985–2023 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Tucker Hill PAC are available in Appendix I (Figure A1.22).




Figure 28. *N*-mixture model estimates of sage-grouse abundance within Tucker Hill Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Warner PAC

The Warner PAC is an area of approximately 135,531 ha (334,901 ac), located within Lake County, and contains 62 known leks. As of 2025, the conservation statuses of these leks are: 33 occupied, 5 occupied-pending, 9 unoccupied, and 15 historic leks.

During 2025, 68 lek surveys were conducted at 29 leks within the Warner PAC; 28 (73.7%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each surveyed lek in the Warner PAC was counted 2.34 times. Estimated sage-grouse abundance in the Warner PAC in 2025 was 4,218 birds (95% CI: 3,321–5,477 birds), up +27.1% from the 3,318 birds estimated in the PAC in 2024 (Figure 29). The Warner PAC population makes up 8.3% of Oregon’s overall sage-grouse population (Table 2).

The short-term population trend (1 oscillation) in the Warner PAC was -14.4% during 2014–2019, the mid-term population trend (3 oscillations) was -46.2% during 1996–2019, and the long-term population trend (4 oscillations) was -65.5% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for the Warner PAC are available in Appendix I (Figure A1.23).




Figure 29. *N*-mixture model estimates of sage-grouse abundance within Warner Priority Area for Conservation in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Areas outside of PACs (Low-Density Habitats)

There are 129 known leks located in Oregon’s low-density habitat areas, outside of PACs. As of 2025, the conservation statuses of these leks are: 24 occupied, 15 occupied-pending, 5 unoccupied-pending, 30 unoccupied, and 55 historic leks.

During 2025, 89 ground surveys and 5 aerial lek surveys were conducted at 51 leks in areas outside of PACs; 29 (74.4%) of the occupied or occupied-pending leks were surveyed one or more times in 2025. On average, each lek surveyed in areas outside of PACs was counted 1.74 times. Estimated sage-grouse abundance in low-density habitat in 2025 was 761 birds (95% CI: 513–1,156 birds), up +15.0% from the 662 birds estimated in the PAC in 2024 (Figure 30).

The short-term population trend (1 oscillation) in areas outside of PACs was -63.1% during 2014–2019, the mid-term population trend (2 oscillations) was -75.6% during 2002–2019, and the long-term population trend (4 oscillations) was -87.1% during 1985–2019 (Table A1.1). Figures illustrating the long-term trends (1980–2024) for areas outside of PACs are available in Appendix I (Figure A1.24).




Figure 30. *N*-mixture model estimates of sage-grouse abundance within low-density habitat areas in the state of Oregon, 2017–2025. The median (black line) and 95% credible interval (gray polygon) are plotted.

Summary and Conclusions

The sage-grouse population in Oregon experienced another large increase in 2025, suggesting populations are still trending upward in Oregon's current population cycle. The most recent population nadir (trough) was evident in 2019 when the statewide population was the lowest estimate recorded during the 46-year period from 1980–2025. Sage-grouse populations exhibit density dependent fluctuations over time (Garton et al. 2011) and population trends (i.e., changes in population size) should be interpreted from complete population cycles using the population nadirs (Coates et al. 2021). In recent decades, sage-grouse populations in Oregon have cycled on an approximately 6–7-year period. The estimated statewide population abundance has followed this pattern over the past 9 years, with 3 years of declining population trends from 2017–19, followed by population increases from 2020–25. During the 1980–2025 timeframe, five nadirs exist in Oregon's statewide sage-grouse population cycles, in years 1985, 1996, 2008, 2013, and 2019. Comparing trends for each of these four previous population nadirs to the most recent nadir, Oregon's sage-grouse populations, statewide, have declined by -32.9% (1985–2019), -20.0% (1996–2019), -11.9% (2008–2019), and -13.8% (2013–2019).

Population increases were expected in 2025 based on analyses of sage-grouse wings collected from hunters following the 2024 hunting season. These wing data indicated sage-grouse production in 2024, as measured by percent juveniles in the harvest and the number of chicks per hen (CPH), was among the highest recorded in Oregon since hunter harvest wing data collection began in 1982. Percent juveniles in the harvest was 55%, above the long-term average of 47% (1993–2023). The number of chicks per hen in 2024 was 2.25 CPH, well above the long-term average of 1.48 CPH (1993–2023).

Previous versions of the annual Oregon Greater Sage-Grouse Population Monitoring Report (2011–2024) presented Oregon's sage-grouse population size estimates and population trends obtained from the breeding population analysis described in Hagen (2011). This methodology was updated in 2025 when ODFW contracted with USGS to develop a new sage-grouse population model (*N*-Mixture model) for Oregon. The updated *N*-Mixture model accounts for imperfect detection of male sage-grouse attending leks and provides unbiased estimates of sage-grouse abundance at large (Oregon range-wide) and smaller (PAC) spatial scales. Importantly, the new model adjusts for presently unknown leks on the landscape by considering those previously unknown leks added to the database annually, 1941–2025. ODFW's Sage-Grouse Population Monitoring Report from 2024 reported a large apparent population increase between 2023 and 2024, largely due to the number of previously unknown leks (35 leks) confirmed active in 2024 and added to the statewide sage-grouse lek database. There have been over 100 new leks discovered in Oregon and added to the statewide lek database over the past 10 years. Under ODFW's previous sage-grouse population model, new leks were mostly unaccounted for, resulting in large apparent population increases following their addition to the lek database.

Under ODFW's updated population model, Oregon's statewide sage-grouse population increased by +17.8% from 2024 to an estimated 50,912 birds in 2025, the highest population estimate recorded in Oregon since 2017. Population increases between 2024 and 2025 occurred in 21 of Oregon's 23 sage-grouse PACs, ranging from +1.5% (Cow Lakes PAC) to +48.1% (Juniper Mountain PAC), with 11 PACs increasing by more than +20%. Decreasing trends occurred in

only 2 PACs, the Baker PAC (-13.3%) and Beatys PAC (-4.5%). Population trends at the statewide level are primarily driven by 8 PACs which make up the majority (>70%) of Oregon's sage-grouse population by abundance: Beatys PAC (14.0%), Steens PAC (10.5%), Louse Canyon PAC (10.4%), Warner PAC (8.3%), Trout Creeks PAC (8.0%), Bully Creek PAC (6.8%), Crowley PAC (6.7%), and Pueblos PAC (6.6%). Although the Beaty PAC declined by -4.5% between 2024 and 2025, large population increases occurred in the Steens PAC (+33.3%), Louse Canyon PAC (+28.0%), Warner PAC (+27%), Trout Creeks PAC (+23.8%), Crowley PAC (+36.7%), and Pueblos PAC (+23.5%), and a moderate increase occurred in the Bully Creek PAC (+11.5%). Population decline in the Beatys PAC in 2025 could be a result of impacts from the Warner Peak wildlife in the summer of 2024.

The sage-grouse population in the Baker PAC was one of two PACs which saw declines between 2024 and 2025. Unlike adjacent populations, the Baker PAC population has consistently declined since 2017. Based on these abundance estimates and the raw lek count data showing little variation in male lek complex attendance over the past 10 years, this population appears to have stabilized at a low level of abundance. Concerningly, sage-grouse population within the Baker PAC has not displayed density dependent fluctuations or responses to changes in annual precipitation patterns over the past decade, like those patterns observed in populations throughout the majority of Oregon's sage-grouse range. The lack of a response to annual conditions by the Baker PAC sage-grouse population may indicate higher probability of extirpation of this local population (Coates et al. 2018). Concern for the persistence of the Baker PAC sage-grouse population has resulted in conversations regarding a potential translocation effort, which would augment this population by bringing birds from adjacent source populations to the Baker PAC, with a goal of increasing genetic diversity within this potentially genetically depressed population. As of November 2025, these discussions are ongoing within the Department and among ODFW's partners in sage-grouse conservation.

ODFW's original sage-grouse population and habitat objectives for Oregon established 2003 as the baseline year in which Oregon's sage-grouse population and habitat metrics should be assessed (Hagen 2005). During the most recent update (2025) to the Oregon Sage-Grouse Conservation Assessment and Strategy (CAAS), the Department updated the statewide sage-grouse population goal and objectives to reflect ODFW's new population model and the best available science regarding unbiased assessment of sage-grouse population trends (Coates et al. 2021). Considering sage-grouse populations in Oregon generally oscillate over 6–12-year cycles, the Department will continue to manage sage-grouse to maintain or enhance their statewide distribution and abundance oscillating around the 2003 spring breeding population level, approximately 53,000 birds, over the next 50 years. As of 2025, the estimated population is -2.7% below the 2003 baseline, and this estimate is the closest Oregon has been to reaching the statewide baseline population since 2006. However, as ODFW's management objectives for Oregon's sage-grouse populations are to maintain stable or increasing population trends statewide and at the PAC-scale, where trends are assessed between nadirs (troughs) of the population cycles, the population trends in many of Oregon's sage-grouse PACs are quite concerning (Appendix I). Only a few PACs have seen stable or increasing population trends in the short-term, mid-term, or long-term. Additionally, Oregon's 2019 statewide sage-grouse population estimate was the lowest recorded during the monitoring timeframe (1980–2024), which still warrants serious concern for the sage-grouse populations in Oregon. The statewide

population has rebounded following the 2019 historic low estimate, and the moderate to large population increases realized over the past 3–4 years in the current population cycle may be adequate to prevent the next trough in the cycle from being a new historic low population estimate for sage-grouse in Oregon, depending on precipitation patterns and wildfire propensity over the next decade. Continued dedication to maintaining high levels of sage-grouse lek survey effort, both ground and aerial, will help ODFW closely monitor the current population trend. Oregon has an excellent framework for sage-grouse conservation through Oregon Administrative Rules (OARs) 660-023-0115 and 635-140-0000–0025, but active habitat management and restoration in core sage-grouse habitat should be accelerated to the greatest extent possible to maintain sage-grouse range at its current extent and support the rebound of Oregon’s sage-grouse populations over the long-term.

ODFW is confident that the sage-grouse population abundance estimates and trends reported in this document are accurate and scientifically supported, as they were informed by ODFW’s best available sage-grouse data, and should be considered the best available science. Standardized lek survey procedures have been in place since 1996, providing reliable, high-quality sage-grouse population data collected in Oregon during the past 30+ years. Additionally, the Department has updated its sage-grouse population analyses to utilize the most current statistical modeling techniques and minimize any potential biases in the population estimates. However, it is important to note that estimating sage-grouse populations from lek surveys is a complicated process, containing multiple assumptions and potential sources of uncertainty, including assumptions regarding the lek attendance rate of male sage-grouse, knowledge of the distribution of leks in an area, bias in the selection of leks to be monitored in a given year, and uncertainty regarding the rate of new lek formation (Beck and Braun 1980, Walsh et al. 2004). The information presented in this report should be interpreted within the context of these assumptions, sources of uncertainty, and potential biases.

References

- Beck, T. D. I., and C. E. Braun. 1980. The strutting ground count: variation, traditionalism, management needs. *Proceedings of the Annual Conference of the Western Association of State Game and Fish Commissioners* 60:558–566.
- Cline, M. L., Vold, S. T., Jackle, G. S. 2025. Greater sage-grouse conservation assessment and strategy for Oregon: a plan to maintain and enhance populations and habitat. Third edition. Oregon Department of Fish and Wildlife, Salem, USA.
- Coates, P. S., B. G. Prochazka, C. Aldridge, M. O'Donnell, D. R. Edmunds, A. P. Monroe, S. Hanser, L. A. Wiechman, and M. P. Chenaille. 2025. Trends and a targeted annual warning system for greater sage-grouse in the western United States (ver. 4.0). U.S. Geological Survey data release.
- Coates, P.S., B. G. Prochazka, M. S. O'Donnell, C. L. Aldridge, D. R. Edmunds, A. P. Monroe, M. A. Ricca, G. T. Wann, S. E. Hanser, L. A. Wiechman, and M. P. Chenaille. 2021. Range-wide greater sage-grouse hierarchical monitoring framework - Implications for defining population boundaries, trend estimation, and a targeted annual warning system:

U.S. Geological Survey Open-File Report 2020-1154.
<https://doi.org/10.3133/ofr20201154>

- Coates, P. S., Prochazka, B. G., Ricca, M. A., Halstead, B. J., Casazza, M. L., Blomberg, E. J., Brussee, B. E., Wiechman, L., Tebbenkamp, J., Gardner, S. C. and K. P. Reese. 2018. The relative importance of intrinsic and extrinsic drivers to population growth vary among local populations of Greater Sage-Grouse: An integrated population modeling approach. *The Auk: Ornithological Advances* 135: 240–261.
- Garton, E. O., J. W. Connelly, J. S. Horne, C. A. Hagen, A. Moser, and M. A. Schroeder. 2011. Greater sage-grouse population dynamics and probability of persistence. Greater sage-grouse: ecology and conservation of a landscape species and its habitats. *Studies in Avian Biology* 38:293–381.
- Hagen, C. A. 2005. Greater sage-grouse conservation assessment and strategy for Oregon: a plan to maintain and enhance populations and habitat. First edition. Oregon Department of Fish and Wildlife, Salem, USA.
- Hagen, C. A. 2011. Greater sage-grouse conservation assessment and strategy for Oregon: a plan to maintain and enhance populations and habitat. Second edition. Oregon Department of Fish and Wildlife, Salem, USA.
- McCaffery, R, J. J. Nowak, P. M. Lukacs. 2016. Improved analysis of lek count data using n-mixture models. *Journal of Wildlife Management* 80:1011–1021.
- Patterson, R. L. 1952. *The sage grouse in Wyoming*. Sage Books Inc., Denver, Colorado, USA.
- United States Department of the Interior Bureau of Land Management Oregon/Washington State Office [US DOI BLM OR/WA]. 2015. Oregon Greater Sage-Grouse Approved Resource Management Plan Amendment.
https://eplanning.blm.gov/public_projects/lup/103348/143725/176961/OR_GRSR_ROD_ARMPA.pdf
- Walsh, D. P., G. C. White, T. E. Remington, and D. C. Bowden. 2004. Evaluation of the lek-count index for greater sage-grouse. *Wildlife Society Bulletin* 32:56–68.

Appendix I – Long-Term Population Trends by PAC

ODFW delineated the breeding habitat, based on lek size and distribution, of approximately 90% of the state’s sage-grouse population, and grouped this area into 20 “Sage-Grouse Core Areas” during 2011–12 (Hagen 2011). During 2022–2023, ODFW worked with our partners to update Oregon’s map of sage-grouse core areas and low-density habitats. The project identified the most productive sage-grouse populations and habitats and the important connectivity habitats among these core habitats. The revision incorporated new data collected since 2011 and utilized the best available science to identify areas that currently provide breeding, brood-rearing and summering, wintering, and migratory habitats for greater sage-grouse in Oregon. In December 2023, the Oregon Fish and Wildlife Commission reviewed and approved the updated habitat map (Figure 1).

Since the initial delineation of these core areas, they have been incorporated into multiple assessments and regulatory documents, including the 2015 U.S. Fish and Wildlife Service “Not-Warranted” decision (USFWS 2015), the BLM Oregon Greater Sage-Grouse Approved Resource Management Plan Amendments (US DOI BLM OR/WA 2015, US DOI BLM 2025), and the Oregon Sage-Grouse Action Plan (SageCon 2015). The term Priority Area of Conservation (PAC) corresponds directly with ODFW’s core areas, and the term Priority Habitat Management Area (PHMA) describes the portions of each core area (PAC) under BLM administration (US DOI BLM 2025). ODFW provides lek survey results to the BLM following the lek survey period and coordinates with BLM during the annual assessment of BLM’s sage-grouse adaptive management triggers. Concurrently with their adoption in various regulatory documents, Oregon’s PACs are a significant focus of ongoing landscape-scale sage-grouse habitat management and conservation.

Habitat management for sage-grouse in Oregon often occurs at the PAC-scale, or at even smaller scales within a PAC. Sage-grouse population abundance estimates from ODFW’s new *N*-Mixture model are unavailable at these finer scales prior to 2017; however, estimates of relative sage-grouse abundance are available through the USGS Rangewide Population Monitoring Framework (Coates et al. 2021, 2025). Note, there is a 1-year lag for these reported trends, as the results from the annual Rangewide Population Monitoring Framework analyses are not available until the following calendar year (i.e., USGS Population Monitoring Report from the 2025 lek survey season will be available in 2026).

Nadir-to-nadir sage-grouse population trends were calculated for each PAC during three time periods, short-term, mid-term, and long-term, using sage-grouse abundance data from Coates et al. 2025 (Ver. 4). Within PACs, the estimated Abundance Indexes for all leks were summed each year, 1980–2024. The short-term trend is defined as the most recent single oscillation of Oregon’s sage-grouse population cycle, the mid-term trend is the most recent 2 or 3 oscillations of the population cycle, and the long-term trend is the most recent 4 or 5 oscillations of Oregon’s sage-grouse population cycle. The specific time periods for population cycle oscillations of each PAC are described below and are viewable in the associated figures, below.

References

- Bureau of Land Management (BLM). 2015. Oregon greater sage-grouse approved resource management plan amendment. US Department of Interior, Bureau of Land Management, Oregon/Washington State Office, Portland, USA. BLM/OR/WA/PL-15/051+1792.
- Coates, P. S., B. G. Prochazka, C. Aldridge, M. O'Donnell, D. R. Edmunds, A. P. Monroe, S. Hanser, L. A. Wiechman, and M. P. Chenaille. 2025. Trends and a targeted annual warning system for greater sage-grouse in the western United States (ver. 4.0). U.S. Geological Survey data release.
- Coates, P.S., B. G. Prochazka, M. S. O'Donnell, C. L. Aldridge, D. R. Edmunds, A. P. Monroe, M. A. Ricca, G. T. Wann, S. E. Hanser, L. A. Wiechman, and M. P. Chenaille. 2021. Range-wide greater sage-grouse hierarchical monitoring framework - Implications for defining population boundaries, trend estimation, and a targeted annual warning system: U.S. Geological Survey Open-File Report 2020-1154.
<https://doi.org/10.3133/ofr20201154>
- Hagen, C. A. 2011. Greater sage-grouse conservation assessment and strategy for Oregon: a plan to maintain and enhance populations and habitat. Second edition. Oregon Department of Fish and Wildlife, Salem, USA.
- Sage-Grouse Conservation Partnership (SageCon). 2015. The Oregon sage-grouse action plan. Oregon Governor's Natural Resource Office, Salem, USA.
- United States Department of the Interior Bureau of Land Management Oregon/Washington State Office [US DOI BLM OR/WA]. 2015. Oregon Greater Sage-Grouse Approved Resource Management Plan Amendment.
https://eplanning.blm.gov/public_projects/lup/103348/143725/176961/OR_GRSR_ROD_ARMPA.pdf
- United States Department of the Interior Bureau of Land Management [US DOI BLM]. 2025. Greater Sage-Grouse Rangewide Planning Record of Decision and Approved Resource Management Plan Amendment for Oregon.
https://eplanning.blm.gov/public_projects/2016719/200502020/20126642/251026622/Oregon%20ROD%20&%20ARMPA%20-%201.15.2025.pdf
- U.S. Fish and Wildlife Service (USFWS). 2015. Endangered and threatened wildlife and plants; 12-month findings for petitions to list the greater sage-grouse (*Centrocercus urophasianus*) as threatened or endangered. Federal Register 80:59857–59942.

Table A1. 1. Sage-grouse population trends within each of Oregon’s Sage-Grouse Priority Areas for Conservation (PACs), 1980–2024.

Priority Area for Conservation (PAC)	Trend		
	Short-term ¹ (1 oscillation, ~2012–2020)	Mid-term ² (2 or 3 oscillations, ~2000–2020)	Long-term ³ (4 or 5 oscillations, ~1985–2020)
Baker	-22.8%	-79.6%	-81.5%
Beatys	-17.4%	-72.7%	-76.5%
Brothers	-58.7%	-53.5%	-80.9%
Bully Creek	-31.1%	-35.8%	-20.9%
Burns	-32.7%	-49.8%	-66.9%
Cow Lakes	-29.6%	-68.7%	-85.4%
Cow Valley	+1.9%	-21.8%	-3.3%
Crowley	-16.9%	-38.1%	-55.5%
Diablo	-42.6%	-60.2%	-71.8%
Drewsey	-31.3%	-43.7%	-73.3%
Dry Valley – Jack Mountain	0.0%	-72.2%	-85.7%
Juniper Mountain	+10.1%	+16.6%	-32.4%
Louse Canyon	-21.2%	-25.0%	-56.8%
North Wagontire	-33.7%	-49.5%	-73.7%
Paulina	-15.4%	-51.7%	-60.3%
Pueblos	-5.8%	-34.1%	-53.2%
Saddle Butte	+55.5%	+40.5%	0.0%
Sheepshead	-24.2%	-48.2%	-65.4%
Soldier Creek	-37.9%	-52.2%	-68.3%
Steens	-10.2%	-10.4%	-45.4%
Trout Creeks	+7.1%	-18.4%	-49.3%
Tucker Hill	-8.4%	+3.6%	+61.1%
Warner	-14.4%	-46.2%	-65.5%
Areas outside of PACs	-63.1%	-75.6%	-87.1%

¹ The short-term trend is defined as the most recent single oscillation of Oregon’s sage-grouse population cycle. Generally, the short-term trend considers the 2012–2020 time period. The trend is calculated among nadirs of the population cycle for each PAC independently, so some variability exists in the defined time periods. Refer to the section below for the specific population nadirs and associated time period oscillations for each PAC.

² The mid-term trend is defined as the most recent 2 or 3 oscillations of Oregon’s sage-grouse population cycle. Generally, the mid-term trend considers the 2000–2020 time period.

³ The long-term trend is defined as the most recent 4 or 5 oscillations of Oregon’s sage-grouse population cycle. Generally, the long-term trend considers the 1985–2020 time period.

Baker PAC

Nadirs of the population cycle in the Baker PAC occurred during the years 1985, 1993, 2001, 2014, and 2023 (Figure A1.1A). The Baker PAC experienced an increasing population trend during the period of 1993–2001, and declining trends during all other periods (Figure A1.1B). The short-term population trend (1 oscillation) was -22.8% during 2014–2023, the mid-term population trend (2 oscillations) was -79.6% during 2001–2023, and the long-term population trend (4 oscillations) was -81.5% during 1985–2023 (Table A1.1).




Figure A1. 1. Sage-grouse population trends in the Baker PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Beatys PAC

Nadirs of the population cycle in the Beatys PAC occurred during the years 1985, 1996, 2002, 2013, and 2019 (Figure A1.2A). The Beatys PAC experienced an increasing population trend during the period of 1996–2002, and declining trends during all other periods (Figure A1.2B). The short-term population trend (1 oscillation) was -17.4% during 2013–2019, the mid-term population trend (2 oscillations) was -72.7% during 2002–2019, and the long-term population trend (4 oscillations) was -76.5% during 1985–2019 (Table A1.1).




Figure A1. 2. Sage-grouse population trends in the Beatys PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Brothers PAC

Nadirs of the population cycle in the Brothers PAC occurred during the years 1985, 1996, 2003, 2009, and 2019 (Figure A1.3A). The Brothers PAC experienced an increasing population trend during the period of 2003–2009, and declining trends during all other periods (Figure A1.3B). The short-term population trend (1 oscillation) was -58.7% during 2009–2019, the mid-term population trend (2 oscillations) was -53.5% during 2003–2019, and the long-term population trend (4 oscillations) was -80.9% during 1985–2019 (Table A1.1).




Figure A1. 3. Sage-grouse population trends in the Brothers PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Bully Creek PAC

Nadirs of the population cycle in the Bully Creek PAC occurred during the years 1985, 1993, 2001, 2008, 2014, and 2020 (Figure A1.4A). The Bully Creek PAC experienced an increasing population trend during the periods of 1993–2001 and 2001–2009, and declining trends during all other periods (Figure A1.4B). The short-term population trend (1 oscillation) was -31.1% during 2014–2020, the mid-term population trend (3 oscillations) was -35.8% during 2001–2020, and the long-term population trend (5 oscillations) was -20.9% during 1985–2020 (Table A1.1).




Figure A1. 4. Sage-grouse population trends in the Bully Creek PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Burns PAC

Nadirs of the population cycle in the Burns PAC occurred during the years 1985, 1996, 2002, 2009, 2014, and 2020 (Figure A1.5A). The Burns PAC experienced an increasing population trend during the period of 1996–2002, and declining trends during all other periods (Figure A1.5B). The short-term population trend (1 oscillation) was -32.7% during 2014–2020, the mid-term population trend (3 oscillations) was -49.8% during 2002–2020, and the long-term population trend (5 oscillations) was -66.9% during 1985–2020 (Table A1.1).




Figure A1. 5. Sage-grouse population trends in the Burns PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Cow Lakes PAC

Nadirs of the population cycle in the Cow Lakes PAC occurred during the years 1985, 1995, 2001, 2008, 2013, and 2022 (Figure A1.6A). The Cow Lakes PAC experienced an increasing population trend during the period of 1995–2001, and declining trends during all other periods (Figure A1.6B). The short-term population trend (1 oscillation) was -29.6% during 2013–2022, the mid-term population trend (3 oscillations) was -68.7% during 2001–2022, and the long-term population trend (5 oscillations) was -85.4% during 1985–2022 (Table A1.1).




Figure A1. 6. Sage-grouse population trends in the Cow Lakes PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Cow Valley PAC

Nadirs of the population cycle in the Cow Valley PAC occurred during the years 1985, 1996, 2001, 2008, 2014, and 2019 (Figure A1.7A) and declining trends during all other periods (Figure A1.7B). The short-term population trend (1 oscillation) was +1.9% during 2014–2019, the mid-term population trend (3 oscillations) was -21.8% during 2001–2019, and the long-term population trend (5 oscillations) was -3.3% during 1985–2019 (Table A1.1).




Figure A1. 7. Sage-grouse population trends in the Cow Valley PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Crowley PAC

Nadirs of the population cycle in the Crowley PAC occurred during the years 1984, 1996, 2002, 2008, 2013, and 2021 (Figure A1.8A). The Crowley PAC experienced an increasing population trend during the period of 2008–2013, and declining trends during all other periods (Figure A1.8B). The short-term population trend (1 oscillation) was -16.9% during 2013–2021, the mid-term population trend (3 oscillations) was -38.1% during 2002–2021, and the long-term population trend (5 oscillations) was -55.5% during 1984–2021 (Table A1.1).




Figure A1. 8. Sage-grouse population trends in the Crowley PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Diablo PAC

Nadirs of the population cycle in the Diablos PAC occurred during the years 1985, 1996, 2002, 2009, 2013, and 2021 (Figure A1.9A). The Diablo PAC experienced an increasing population trend during the period of 1996–2002, and declining trends during all other periods (Figure A1.9B). The short-term population trend (1 oscillation) was -42.6% during 2013–2021, the mid-term population trend (3 oscillations) was -60.2% during 2002–2021, and the long-term population trend (5 oscillations) was -71.8% during 1985–2021 (Table A1.1).




Figure A1. 9. Sage-grouse population trends in the Diablo PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Drewsey PAC

Nadirs of the population cycle in the Drewsey PAC occurred during the years 1985, 1996, 2002, 2009, 2013, and 2019 (Figure A1.10A). The Drewsey PAC experienced an increasing population trend during the period of 2002–2009, and declining trends during all other periods (Figure A1.10B). The short-term population trend (1 oscillation) was -31.3% during 2013–2019, the mid-term population trend (3 oscillations) was -43.7% during 2002–2019, and the long-term population trend (5 oscillations) was -73.3% during 1985–2019 (Table A1.1).




Figure A1. 10. Sage-grouse population trends in the Drewsey PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Dry Valley – Jack Mountain PAC

Nadirs of the population cycle in the Dry Valley – Jack Mountain PAC occurred during years 1966, 1974, 1984, 1996, 2002, 2009, 2015, and 2019 (Figure A1.11A). The Dry Valley – Jack Mountain PAC experienced an increasing population trend during the period of 1996–2002, and declining trends during all other periods (Figure A1.11B). The short-term population trend (1 oscillation) was stable (0.0%) during 2015–2019, the mid-term population trend (3 oscillations) was -71.2% during 2002–2019, and the long-term population trend (5 oscillations) was -85.7% during 1984–2019 (Table A1.1).




Figure A1. 11. Sage-grouse population trends in the Dry Valley – Jack Mountain PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Juniper Mountain PAC

Nadirs of the population cycle in the Juniper Mountain PAC occurred during the years 1984, 1996, 2009, 2013, and 2019 (Figure A1.12A). The Juniper Mountain PAC experienced an increasing population trend during the periods of 1996–2009 and 2013–2019, and declining trends during all other periods (Figure A1.12B). The short-term population trend (1 oscillation) was +10.1% during 2013–2019, the mid-term population trend (3 oscillations) was +16.6% during 1996–2019, and the long-term population trend (4 oscillations) was -32.4% during 1984–2019 (Table A1.1).




Figure A1. 12. Sage-grouse population trends in the Juniper Mountain PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Louse Canyon PAC

Nadirs of the population cycle in the Louse Canyon PAC occurred during the years 1985, 1996, 2014, and 2019 (Figure A1.13A). The Louse Canyon PAC experienced declining trends during all time periods (Figure A1.13B). The short-term population trend (1 oscillation) was -21.2% during 2014–2019, the mid-term population trend (2 oscillations) was -25.0% during 1996–2019, and the long-term population trend (3 oscillations) was -56.8% during 1985–2019 (Table A1.1). Note, unlike most PACs, the abundance data do not suggest the Louse Canyon PAC experienced a population nadir during the 2000–2010 period.




Figure A1. 13. Sage-grouse population trends in the Louse Canyon PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

North Wagontire PAC

Nadirs of the population cycle in the North Wagontire PAC occurred during the years 1993, 2002, 2013, and 2023 (Figure A1.14A). The North Wagontire PAC experienced declining population trends during all time periods (Figure A1.14B). The short-term population trend (1 oscillation) was -33.7% during 2013–2023, the mid-term population trend (2 oscillations) was -49.5% during 2002–2023, and the long-term population trend (4 oscillations) was -73.7% during 1985–2023 (Table A1.1).




Figure A1. 14. Sage-grouse population trends in the North Wagontire PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Paulina PAC

Nadirs of the population cycle in the Paulina PAC occurred during the years 1985, 1996, 2002, 2009, 2013, and 2023 (Figure A1.15A). The Paulina PAC experienced an increasing population trend during the period of 2009–2013, and declining population trends during all time periods (Figure A1.15B). The short-term population trend (1 oscillation) was -15.4% during 2013–2023, the mid-term population trend (3 oscillations) was -51.7% during 2002–2023, and the long-term population trend (5 oscillations) was -60.3% during 1985–2023 (Table A1.1).




Figure A1. 15. Sage-grouse population trends in the Paulina PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Pueblos PAC

Nadirs of the population cycle in the Pueblos PAC occurred during the years 1985, 1996, 2009, 2013, 2019, and 2022 (Figure A1.16A). The Pueblos PAC experienced an increasing population trend during the period of 1996–2009, and declining trends during all other periods (Figure A1.16B). The short-term population trend was -5.8% during 2019–2022 (1 oscillation) and -31.8% during 2013–2022 (2 oscillations). The mid-term population trend (3 oscillations) was -31.4% during 2009–2022, and the long-term population trend (5 oscillations) was -53.2% during 1985–2022 (Table A1.1).




Figure A1. 16. Sage-grouse population trends in the Pueblos PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Saddle Butte PAC

Nadirs of the population cycle in the Saddle Butte PAC occurred during the years 1985, 1996, 2002, 2009, 2013, and 2020 (Figure A1.17A). The Saddle Butte PAC experienced an increasing population trend during the periods of 1996–2002 and 2013–2020, and declining trends during all other periods (Figure A1.17B). The short-term population trend (1 oscillation) was +55.5% during 2013–2020, the mid-term population trend (3 oscillations) was +40.5% during 2002–2020, and the long-term population trend (5 oscillations) was stable (+0.0%) during 1985–2020 (Table A1.1).




Figure A1. 17. Sage-grouse population trends in the Saddle Butte PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Sheepshead PAC

Nadirs of the population cycle in the Sheepshead PAC occurred during the years 1985, 1996, 2002, 2009, 2013, and 2021 (Figure A1.18A). The Sheepshead PAC experienced an increasing population trend during the period of 1996–2002, and declining trends during all other periods (Figure A1.18B). The short-term population trend (1 oscillation) was -24.2% during 2013–2021, the mid-term population trend (3 oscillations) was -48.2% during 2002–2021, and the long-term population trend (5 oscillations) was -65.4% during 1985–2021 (Table A1.1).




Figure A1. 18. Sage-grouse population trends in the Sheepshead PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Soldier Creek PAC

Nadirs of the population cycle in the Soldier Creek PAC occurred during the years 1985, 1996, 2008, 2013, and 2020 (Figure A1.19A). The Soldier Creek PAC experienced declining population trends during all time periods (Figure A1.19B). The short-term population trend (1 oscillation) was -37.9% during 2013–2020, the mid-term population trend (3 oscillations) was -52.2% during 1996–2020, and the long-term population trend (4 oscillations) was -75.2% during 1985–2020 (Table A1.1).




Figure A1. 19. Sage-grouse population trends in the Soldier Creek PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Steens PAC

Nadirs of the population cycle in the Steens PAC occurred during the years 1985, 1996, 2008, 2013, and 2020 (Figure A1.20A). The Steens PAC experienced an increasing population trend during the period of 1996–2008, and declining trends during all other periods (Figure A1.20B). The short-term population trend (1 oscillation) was -10.2% during 2013–2020, the mid-term population trend (3 oscillations) was -10.4% during 1996–2020, and the long-term population trend (4 oscillations) was -45.4% during 1985–2020 (Table A1.1).




Figure A1. 20. Sage-grouse population trends in the Steens PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values > 1.0 indicates a positive population trend during the time period.

Trout Creeks PAC

Nadirs of the population cycle in the Trout Creeks PAC occurred during the years 1985, 1996, 2002, 2008, 2013, and 2021 (Figure A1.21A). The Trout Creeks PAC experienced an increasing population trend during the period of 1996–2002 and 2013–2021, and declining trends during all other periods (Figure A1.21B). The short-term population trend (1 oscillation) was +7.1% during 2013–2021, the mid-term population trend (3 oscillations) was -18.4% during 2002–2021, and the long-term population trend (5 oscillations) was -49.3% during 1985–2021 (Table A1.1).




Figure A1. 21. Sage-grouse population trends in the Trout Creeks PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Tucker Hill PAC

Nadirs of the population cycle in the Tucker Hill PAC occurred during the years 1984, 1997, 2001, 2009, 2013, and 2023 (Figure A1.22A). The Tucker Hill PAC experienced an increasing population trend during the periods of 1984–1997 and 2001–2009, and declining trends during all other periods (Figure A1.22B). The short-term population trend (1 oscillation) was -8.4% during 2013–2023, the mid-term population trend (3 oscillations) was +3.6% during 2001–2023, and the long-term population trend (5 oscillations) was +61.1% during 1985–2023 (Table A1.1).




Figure A1. 22. Sage-grouse population trends in the Tucker Hill PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Warner PAC

Nadirs of the population cycle in the Warner PAC occurred during the years 1985, 1996, 2009, 2014, and 2019 (Figure A1.23A). The Warner PAC experienced declining population trends during all time periods (Figure A1.23B). The short-term population trend (1 oscillation) was -14.4% during 2014–2019, the mid-term population trend (3 oscillations) was -46.2% during 1996–2019, and the long-term population trend (4 oscillations) was -65.5% during 1985–2019 (Table A1.1).




Figure A1. 23. Sage-grouse population trends in the Warner PAC. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Areas outside of PACs

Nadirs of the population cycle in areas outside of PACs occurred during the years 1985, 1996, 2002, 2009, and 2019 (Figure A1.24A). Areas outside of Oregon’s PACs experienced declining population trends during all time periods (Figure A1.24B). The short-term population trend (1 oscillation) was -63.1% during 2014–2019, the mid-term population trend (2 oscillations) was -75.6% during 2002–2019, and the long-term population trend (4 oscillations) was -87.1% during 1985–2019 (Table A1.1). Notably, those areas outside of PACs have realized greater declines in population trend in both the short-term and long-term than areas within PACs.




Figure A1. 24. Sage-grouse population trends in areas outside of Oregon’s PACs. Panel A shows the index of sage-grouse abundance through time, 1980–2024. Panel B shows the proportional change in the index of sage-grouse abundance from nadir-to-nadir. Value 1.0 indicates a stable population trend during the time period, values < 1.0 indicates a negative population trend during the time period, and values >1.0 indicates a positive population trend during the time period.

Appendix II – 2025 ODFW Adopt-A-Lek Program Report



2025 Volunteer Field Report

Southeast Oregon Greater Sage-Grouse Adopt-A-Lek Program

*Oregon Department of Fish and Wildlife
Prepared by Damian Fagan*

Summary

This year marked the 20th year of ODFW’s Adopt-A-Lek (AAL) volunteer program. The program provides an opportunity for citizen scientists from Oregon, Washington, and Idaho to count greater sage-grouse (*Centrocercus urophasianus*) leks in remote portions of Malheur County. Data collected by the volunteers contribute to ODFW’s annual population estimate for the species, which is essential to monitoring the health of the sage-grouse populations in Oregon.

This year, 55 AAL volunteers signed up to participate in the counts. Unfortunately, several had to cancel their assignments due to weather and road conditions. It was a challenging year to access areas due to the intermittent storms, muddy roads, and snow.

This year we had new volunteers from Port Townsend and Vancouver, WA; from Eugene, Ontario, Lakeview, Hermiston, Salem, and Long Creek, OR; and Fruitland, ID. Thanks for joining the program!

The 2025 lek counting effort in Oregon continued to focus on key habitat areas referred to as Priority Areas for Conservation (PACs). Although these PACs contain high priority trend leks, very few were counted three times between March 15 and April 30, with 7–10 days spaced

between each count morning. Again, weather and road access had an impact on areas not getting surveyed. Therefore, the value of each count morning cannot be understated.

A total of 136 leks were assigned to the AAL program this year and 121 of them were counted at least once. Fifteen leks were not counted due to weather, difficult access, or road conditions. Many of these fifteen leks are in remote areas with very rough access and rarely get counted.

This year, 295 dawn counts were conducted on 121 leks which resulted in a total of 2075 individual birds counted. Of course, some of these numbers reflect birds counted on multiple surveys. Sixty-two leks were active (having 1 or more displaying males) and roughly 51 percent of the leks had some level of activity (one lek had activity but the birds were present were hens). Of the 62 leks that had activity, 42% of the leks had 1–10 male grouse present, 29% had 11–20 males present, and 29% had more than 20 males present. The highest count of males on a lek was 64 birds, with a total of 74 birds counted on that lek.

Additional grouse were observed but absent during the dawn lek counts, having been observed either during mid-day lek checks or during travel to or away from the lek areas. Several suspected leks were checked and two potential new leks were located.

Volunteers contributed over 917 hours at an estimated value of over \$32,614 (not all volunteer hour forms were returned) and this does not include amounts that were donated back to the program from travel reimbursements.

Several interesting notes from this season:

- Several leks which have not had displaying males for several years had activity this year.
- Two possible new leks were discovered with activity.
- The total highest male counts for all the leks in 2025 was 1049. In 2024 it was 809.
- Of all the leks that were active, 29% had over 21 males present on a given count. In 2024, that number was 24% had over 21 males present on a given count.
- Several leks, which are near one another, had birds moving between the leks, as revealed from the dawn counts. In 2026, it would be good to have nearby leks counted on the same count morning to determine if birds are moving between the leks.
- Western Rivers Conservancy, ODFW, Wilkinson Ranch, and the AAL program hosted an Oregon Field Guide film crew this spring at the Disaster Peak Ranch. Not sure on the release date for the segment.

Even though weather and road conditions were at times extremely challenging, AAL volunteers persevered and documented new observation points, lek activity, and lek access. Many volunteers made special efforts to write detailed comments and observations, share photos, and make suggestions to improve the AAL program which is always appreciated. And a special shout-out to April Wilkinson for fielding calls and emails on the weather and road conditions in the McDermitt area, and to Western Rivers Conservancy for use of the Disaster Peak Ranch's bunkhouse. ODFW seasonal staff Chip Dale and Russ Morgan also provided information to AAL volunteers regarding road conditions.

The AAL count data adds significantly to the statewide Sage-Grouse population database used by state biologists and federal land managers to manage this sagebrush-obligate species. Many thanks to the Oregon Wildlife Foundation and the Bureau of Land Management for providing funding and assistance and a huge THANK YOU to all the AAL volunteers that contributed to

this program and who share a deep commitment to the conservation and preservation of this charismatic species.

2025 Volunteer Statistics

Note: this is not a year-to-year comparison of the exact same leks. Each year varies in terms of the number of leks assigned to the AAL program and how many leks were surveyed by the program during the season.

- 55 volunteers volunteered to count leks; 52 made it out to the field.
- 121 individual leks were counted (compared to 100 in 2024, 75 in 2023, 106 in 2022, 100 in 2021, 80 in 2020, 91 in 2019, 71 in 2018, 76 in 2017, 63 in 2016, 46 in 2015, 63 in 2014).
- 55 individual leks were counted at least 3 times (compared to 36 in 2024, 13 in 2023, 54 in 2022, 45 in 2022, 35 in 2020, 45 in 2019, 18 in 2018, 17 in 2017, 16 in 2016, 12 in 2015, 0 in 2014). One lek was counted 5 times.
- 295 count mornings were conducted (compared to 207 in 2024, 137 in 2023, 267 in 2022, 243 in 2021, 185 in 2020, 236 in 2019, 171 in 2018, 171 in 2017, 116 in 2016, 89 in 2015, 67 in 2014).
- 51% of the leks counted were active - had birds displaying during the count morning (compared to 52% in 2024, 53% in 2023, 49% in 2022, 45% in 2021, 44% in 2020, 53% in 2019, 51% in 2018, 53% in 2017, 52% in 2016, 72% in 2015, 56% in 2014). One lek that was not considered active had several hens but no displaying males present.
- 59 leks were not active (no birds present) on any count morning (compared to 48 in 2024, 35 in 2023, 56 in 2022, 55 in 2021, 55 in 2020, 44 in 2019 and 35 in 2018)
- 62 leks were active (at least 1 male displaying) on a count morning (compared to 51 in 2024, 40 in 2023, 50 in 2022, 45 in 2022, 35 in 2020, 47 in 2019, 36 in 2018, 40 in 2017, 33 in 2016, 33 in 2015, 35 in 2014). One lek had activity but the birds were listed as “unclassified.”
- 2075 birds were counted on all the dawn counts (compared to 1248 in 2024, 578 in 2023, 942 in 2022, 937 in 2021, 419 in 2020, 962 in 2019, 975 in 2017, 1052 in 2016, 871 in 2015, 453 in 2014). Of this number, the total of high counts for males at a given lek was 1049.
- 0 leks were checked but not counted (compared to 0 in 2024, 0 in 2023, 0 in 2022, 11 in 2021, 0 in 2020, 9 in 2019, 5 in 2018, 2 in 2017, 10 in 2016, 14 in 2015, 34 in 2014).
- 42% of the active leks counted had 1-10 males (compared to 24% in 2024, 55% in 2023, 66% in 2022, 73% in 2021, 70% in 2020, 64% in 2019, 58% in 2018, 55% in 2017, 42% in 2016, 48% in 2015, and 44% in 2014).
- 29% of the active leks counted had 11-20 males (compared to 28% in 2024, 38% in 2023, 20% in 2022, 13% in 2021, 12% in 2020, 19% in 2019, 33% in 2018, 18% in 2017, 21% in 2016, 18% in 2015, and 38% in 2014).
- 29% of the active leks counted had 21 or more males (compared to 24% in 2024, 8% in 2023, 14% in 2022, 13% in 2021, 15% in 2020, 15% in 2019, 14% in 2018, 28% in 2017, 36% in 2016, 33% in 2015, and 15% in 2014).
- The largest lek count had 64 males (74 total) (compared to 49 in 2024, 34 in 2023, 31 in 2022, 43 in 2021, 27 in 2020, 49 in 2019, 40 in 2018, 54 in 2017, 60 in 2016, 41 in 2015, and 37 in 2014).

Funding and support for the 2025 Volunteer Program was provided by the Oregon Department of Fish and Wildlife, Oregon Wildlife Foundation, and Bureau of Land Management. In-kind support was provided by Western Rivers Conservancy.



Gallery of Images










Photo Credits

Cover: Greater Sage-Grouse (Raj Ramamoorthy), Gallery of Images: Sunrise Survey (Becky Jarnes), Sagebrush Sunrise (Laura Brockinton), Mud Bath (Jon Hansen), Curious Cow (Jon Hansen), Road Ruts (Karl Findling), More Road Ruts (Damian Fagan), Josh and Raven Counting (Damian Fagan), McDermitt Bighorn Herd (Damian Fagan).

Appendix III – 2025 Aerial Survey Summary

Oregon Department of Fish and Wildlife (ODFW) conducts annual aerial lek searches, with the goals of 1) locating previously undocumented leks, 2) documenting shifts in sage-grouse breeding distribution, and 3) monitoring activity of leks which are inaccessible from the ground. ODFW contracted aerial services for lek searches by helicopter during spring 2025. Funding for 2025 aerial lek surveys was provided by the BLM through the current ODFW/BLM Cooperative Funding Agreement.

Aerial Lek Searches Overview

Aerial lek searches are conducted by helicopter from ½ hour before to 2 hours after sunrise, following fixed transects separated by ¼ – ½ mile. During searches, the helicopter maintains an altitude of 50–150 feet above ground level and a speed of approximately 60 mph. Helicopter searches and surveys are primarily directed towards the assessment of lek occupancy, as lekking sage-grouse are sensitive to aerial predators, and thus often limit their display behavior in presence of a helicopter. For this reason, following the discovery of previously unknown leks, ground observation of a site is required to confirm lek occupancy and attendance. Counts conducted from a helicopter are generally not used to estimate population trend in an analysis area. Rather, these counts typically function as presence-absence assessments of lek activity. The exception to this rule is when male counts recorded from a helicopter survey are greater than male counts recorded by a ground observer. In these cases, the aerial counts are used to assess both population abundance and trends.

2025 Aerial Survey

During early April 2025, ODFW contracted JL Aviation, Inc. to conduct 43 hours of helicopter lek searches in the Steens PAC and the low-density sage-grouse habitat north of the Sheepshead PAC in Harney County and Malheur County. Approximately 1,600 miles of transects were flown over the course of 7 days, utilizing 2 helicopters (Figure A3.1). The aerial survey documented over 19 locations where sage-grouse have not previously been known to display. Notably, a few of these potential new leks in the Steens PAC are among the highest elevation leks in Oregon, with birds displaying at several locations above 7,000 feet. Of these 19 potential new lek locations, 7 leks were confirmed active by a ground observer in 2025 following the aerial survey, 6 in the Steens PAC and 1 lek in the low-density habitat north of the Sheepshead PAC. These 7 new leks will be confirmed on the ground again in spring 2026 before they are added to the statewide database. The additional potential new lek locations in the Steens PAC and surrounding low-density habitat will need to be verified on the ground during the next several breeding seasons before they can be confirmed as new lek locations and added to the database.

2024 Aerial Survey

During early April 2024, ODFW contracted JL Aviation, Inc. to conduct 64 hours of helicopter lek searches in the Pueblos and Steens PACs and surrounding low-density sage-grouse habitat in Harney County. Approximately 2,500 miles of transects were flown over the course of 8 days, utilizing 2–3 helicopters. The aerial survey documented 16 locations where sage-grouse have not previously been known to display. Of the 16 potential new lek locations, 9 leks were confirmed

active by a ground observer in 2024 following the aerial survey, 4 in the Steens PAC and 5 in the Pueblos PAC. During the 2025 breeding season, in addition to confirming these 9 new leks on the ground, 4 previously unknown leks were checked and confirmed, 3 new leks in the Steens PAC and 1 new lek in the Pueblos PAC. The combined maximum count at these four new leks confirmed in 2025 was 43 males. Three potential new leks discovered during the 2024 aerial survey in the Steens and Pueblos PACs remain to be checked on the ground during the spring 2026 sage-grouse breeding season.

2023 Aerial Survey

During late March through mid-April 2023, ODFW and Hart Mountain National Antelope Refuge contracted JL Aviation, Inc. to conduct over 100 hours of helicopter lek searches in the Beatys PAC (on-refuge and off-refuge) and surrounding low-density sage-grouse habitat in Lake County and a small portion of Harney County. This effort resulted in the discovery of 14 previously unknown leks (10 on Hart Mountain National Antelope Refuge, and 4 off-refuge) which were located by air and confirmed on the ground during 2023–2025. There are no potential new leks discovered during the 2023 aerial survey in the Beatys PAC which remain to be checked on the ground.

2022 Aerial Survey

During mid-March through early April 2022, ODFW conducted helicopter lek searches in the Louse Canyon PAC and a small portion of the Trout Creeks PAC. This effort resulted in the discovery of 12 previously unknown leks that were located by air and confirmed on the ground during subsequent breeding seasons, 2022–2025; 8 new leks in the Louse Canyon PAC and 4 new leks in the Trout Creeks PAC. Two previously unknown leks discovered during the 2022 aerial survey were confirmed on the ground during the 2025 lek survey season and added to the statewide lek database, one lek in the Louse Canyon PAC and one lek in the Trout Creeks PAC. The combined maximum count at these two new leks confirmed in 2025 was 10 males. Five potential new leks discovered during the 2022 aerial survey in the Louse Canyon PAC remain to be checked on the ground during the spring 2026 sage-grouse breeding season.

2021 Aerial Survey

During late-March 2021, ODFW conducted helicopter lek searches in the Soldier Creek PAC. This effort resulted in the discovery of ten previously unknown leks that were located by air and confirmed on the ground during subsequent spring breeding seasons, 2021–2024. One potential new lek discovered during the 2021 aerial survey in the Soldier Creek PAC remains to be checked on the ground during the spring 2026 sage-grouse breeding season.




Figure A3. 1. Greater sage-grouse aerial lek search transects in southeastern Oregon, including portions of the Steens PAC and adjacent low-density habitat, and the low-density habitat north of the Sheepshead PAC, 2025.

Appendix IV – 2025 Sage-Grouse Wing-Bee Report (2024 Hunting Season Data)



Oregon Sage-Grouse Wing Analyses, 2024

Annual Report

Skyler Vold, Sage-Grouse Conservation Coordinator

Executive Summary

Oregon Department of Fish and Wildlife authorized 670 greater sage-grouse (*Centrocercus urophasianus*; hereafter: sage-grouse) permits across 9 Wildlife Management Units during September 2024. Following the 2024 hunting season, 310 sage-grouse wings were received from hunters. This represents the largest wing collection since 2016, when 845 permits were authorized. Production in 2024, as measured by percent juveniles in the harvest and the number of chicks per hen (CPH), was among the highest recorded in Oregon since hunter harvest sage-grouse wing data collection began in 1982. Percent juveniles in the harvest was 55%, above the long-term average of 47% (1993–2023). The number of chicks per hen in 2024 was 2.25 CPH, lower than the record high 2023 production value of 2.50 CPH, but well above the long-term average of 1.48 CPH (1993–2023). Apparent nest success in 2024, based on retention of the primary 9 (P9) feather of harvested females, was below average [P9 nest success: 2024 = 36%; 1993–2023 average = 42%]. Production data collected from hunter-harvested wings in 2024 suggests that sage-grouse populations should be stable or increase in 2025, as our indices of annual production, chicks per hen and proportion juveniles in the harvest, have been above average for 3 consecutive years, 2022–2024.

Overview

In 2024, the sage-grouse hunting season in Oregon was by permit for nine days (7–15 Sept), with a daily and seasonal bag limit of two birds. The hunting season length during 2005–2024 was nine days, compared to five days during 1995–2004, and two days in 1993 and 1994. There have not been any changes to the daily or seasonal bag limit since 1993 (Braun et al. 2015; Table A4.1).

Plumage characteristics of wings are used to assess age and sex of numerous game bird species. By assessing plumage characteristics from hunter-harvested wings, demographic parameters (e.g. age structure, sex ratio, and nest success) can be estimated for sage-grouse populations. Sage-grouse wings have been analyzed to gather information regarding population structure and demography in Oregon since 1982. However, methods used to determine age and

sex from wing characteristics were refined in 1993. Due to this change in methodology, all long-term average rates are calculated only for the 1993–2024 period. As in previous years, all hunters who were successful in the controlled sage-grouse hunt drawing were provided envelopes for the return of sage-grouse wings to Oregon Department of Fish and Wildlife (ODFW). Sage-grouse wings collected during the 2024 hunting season were processed by ODFW and U.S. Fish and Wildlife Service biologists at the annual Wing Bee held on 12 February 2025.

Following the 2024 hunting season, 310 hunter-harvested wings were received from the ten Wildlife Management Units or sub-units (WMUs or sub-WMUs) where permits were allocated (Table A4.2, Figure A4.1). This represents an increase in wing collection from the previous year (2023 = 247 wings), and is the highest wing collection since 2016. Wing collection remains below the 31-year (1993–2023) average of 450 wings (Table A4.3), due to reductions in the number of controlled hunt permits authorized over the past decade. Over 75% of wings ($n = 236$) were received from four WMUs [Beatys Butte, Beulah, Warner, and Whitehorse]; 2 wings were received from the Owyhee WMU, and no wings were received from the Silvies WMU (Table 4). The Falls Wildfire of July–August 2024 likely impacted hunter participation in the Silvies WMU during the 2024 sage-grouse hunting season. Due to concerns about decreasing population trends and the impacts of large wildfires, no permits have been offered in the Sumpter and Lookout Mountain WMUs (WMUs 51 and 64), South Wagontire Sub-WMU (Sub-WMU 73B), and Juniper WMU (WMU 71) since 2014, 2016, and 2017, respectively (Figure A4.1). In 2019, permits were once again offered in that part of the Whitehorse WMU west of Highway 95 and south of the Whitehorse Ranch Road (Trout Creek Mountains, Sub-WMU 68A; Figure A4.1). No permits were offered in this area during 2012–2018 due to ongoing research investigating the impacts of the Holloway Fire on sage-grouse in the Trout Creek Mountains. In 2019, initial research questions had been addressed and these populations had sufficiently rebounded to allow a conservative offering of 25 sage-grouse permits in the subunit. This tag draw was conducted separately from the remainder of the Whitehorse WMU (E. Whitehorse, Sub-WMU 68B; Figure A4.1), to maintain a conservative harvest level in the Trout Creek Mountains sub-unit. In 2020 and 2021, 30 permits were offered in the Trout Creek Mountains sub-unit, followed by an additional 5 permits (35 total) in 2022 and 2023. In 2024, 40 permits were offered in the Trout Creek Mountains sub-unit (Table A4.1).

Table A4. 1. Sage-grouse hunting season dates, lengths, daily and season bag limits, and permits authorized, Oregon, 1993–2024.

Year	Season Date	No. Days	Daily Bag	Season Limit	Permits Authorized
1993	18–19 Sept	2	2	2	865
1994	17–18 Sept	2	2	2	<i>No Data</i>
1995	9–13 Sept	5	2	2	1040
1996	7–11 Sept	5	2	2	1140
1997	6–10 Sept	5	2	2	1265
1998	12–16 Sept	5	2	2	1265
1999	11–15 Sept	5	2	2	1265
2000	9–13 Sept	5	2	2	1265
2001	8–12 Sept	5	2	2	1265
2002	7–11 Sept	5	2	2	1250
2003	6–10 Sept	5	2	2	1275
2004	11–15 Sept	5	2	2	1275
2005	10–18 Sept	9	2	2	1300
2006	9–17 Sept	9	2	2	1300
2007	8–16 Sept	9	2	2	1175
2008	6–14 Sept	9	2	2	1175
2009	12–20 Sept	9	2	2	1150
2010	11–19 Sept	9	2	2	1150
2011	10–18 Sept	9	2	2	1130
2012	8–16 Sept	9	2	2	885
2013	7–15 Sept	9	2	2	870
2014	6–14 Sept	9	2	2	845
2015	12–20 Sept	9	2	2	845
2016	10–18 Sept	9	2	2	845
2017	9–17 Sept	9	2	2	800
2018	8–16 Sept	9	2	2	740
2019	7–15 Sept	9	2	2	645
2020	12–20 Sept	9	2	2	630
2021	11–19 Sept	9	2	2	635
2022	10–18 Sept	9	2	2	640
2023	9–17 Sept	9	2	2	640
2024	7–15 Sept	9	2	2	670

Table A4. 2. Oregon wildlife management units or sub-units with permitted sage-grouse harvest, 2024.

WMU #	WMU Name	2024 Permits
65	Beulah	125
66	Malheur River	100
67	Owyhee	50
68A	Trout Creek Mtns.	40

68B	E. Whitehorse	100
69	Steens Mtn.	50
70	Beatys Butte	90
72	Silvies	20
73 ^a	Wagontire	25
74	Warner	70

^aUnit partially closed to hunting in 2024.




Figure A4. 1. Oregon wildlife management units with permitted greater sage-grouse hunting and the distribution of greater sage-grouse in Oregon, 2024.

Age and Sex Composition

Sage-grouse wings were classified by age (juvenile = hatch year; yearling = second year; adult = after hatch year), based on characteristics of the outer primaries (P7–P10), first secondary, tertials, and wing coverts (Braun and Schroeder 2015). In areas where most of the breeding occurs in March and early April, including Oregon, few yearling males will be identifiable in the harvest due to molt progression (Braun and Schroeder 2015). Additionally, if non-nesting or early nesting yearling females complete their wing molt before harvest, there is no reliable way to differentiate these birds from after second year adult females (Braun and Schroeder 2015). Therefore, in Oregon, after hatch year birds are classified as adults unless they can be definitively identified as yearlings by the presence of juvenile primary 9 (P9) and/or primary 10 (P10) feathers (Braun et al. 2015).

Sex classification was assigned based on the length of the P9 and/or P10 feather, depending on the condition of the wing (Braun and Schroeder 2015). Relatively few yearlings are identified in the harvest in Oregon (31-year average = 6% of harvest); 7% of the harvested birds were classified as yearlings in 2024. Production in 2024, as measured by percent juveniles in the harvest, was among the highest recorded in Oregon since hunter harvest sage-grouse wing data collection began in 1982. Overall, percent of juveniles in the harvest was 55%, representing a decrease from the record high in 2023 (61%), but above the long-term (31-year) average of 47%. Within individual WMUs, the percent of juveniles in the harvest was variable, likely due to small sample sizes (Figure A4.2). The number of wings received for individual WMUs ranged from 0 to 64 (Table A4.4). The greatest proportion of juveniles in the harvest was in the North Wagontire sub-WMU (64%; 7 of 11 wings), followed by the Beulah WMU (62%; 28 of 45 wings), and the Warner WMU (60%; 29 of 48 wings; Table 4; Figure A4.2). The sex ratio of juveniles in the 2024 harvest was skewed heavily female (38 males to 62 females), compared to the long-term average juvenile sex ratio (46:54; Table A4.5).




Figure A4. 2. Nest success, percent juveniles in the harvest, and chicks per hen by WMU where sage-grouse harvest occurred, Oregon, 2024. Missing data in the Beulah and Owyhee WMUs reflects apparent nest success of 0 for the hens harvested in these WMUs in 2024. This is an issue of sample size, as both WMUs recorded chicks in the harvest. No wings were received from the Silvies WMU from the 2024 hunting season.

Table A4. 3. Sex composition by age class, and age composition of harvested sage-grouse, all wildlife management units open to harvest, Oregon, 1993–2024.

Year	N	Juvenile			Yearling			Adult		
		M (%)	F (%)	% Harvest	M (%)	F (%)	% Harvest	M (%)	F (%)	% Harvest
1993	439	51	49	47	26	74	4	40	60	49
1994	764	47	53	43	12	88	7	32	68	50
1995	456	42	58	36	5	95	5	32	68	60
1996	493	42	58	51	4	96	5	31	69	44
1997	586	47	53	54	16	84	4	39	61	39
1998	466	48	52	49	6	94	4	39	61	47
1999	671	46	54	56	14	86	5	41	59	39
2000	592	46	54	44	22	78	8	47	53	48
2001	670	50	50	54	10	90	7	44	56	38
2002	648	51	49	58	9	91	7	46	54	36
2003	655	46	54	48	12	88	5	47	53	47
2004	778	45	55	52	9	91	6	40	60	42
2005	829	46	54	45	5	95	5	46	54	50
2006	669	46	54	47	30	70	5	49	51	48
2007	485	44	56	28	10	90	6	38	62	66
2008	443	49	51	54	0	100	4	30	70	42
2009	493	47	53	57	0	100	5	49	51	38
2010	463	43	57	48	4	96	5	36	64	47
2011	422	43	57	42	10	90	5	48	52	53
2012	321	40	60	29	30	70	14	49	51	57
2013	254	50	50	58	11	89	7	36	64	35
2014	264	38	62	31	6	94	6	42	58	63
2015	290	43	57	58	14	86	2	40	60	40
2016	331	54	46	46	32	68	8	45	55	47
2017	270	42	58	39	5	95	7	37	63	53
2018	255	36	64	42	15	85	5	49	51	53
2019	145	44	56	48	0	100	5	38	62	47
2020	175	44	56	43	7	93	8	37	63	49
2021	150	44	56	43	7	93	8	35	65	53
2022	228	37	63	51	27	73	7	48	52	42
2023	247	57	43	61	0	100	4	43	57	35
<i>31-yr Avg.</i>	<i>450</i>	<i>46</i>	<i>54</i>	<i>47</i>	<i>12</i>	<i>88</i>	<i>6</i>	<i>41</i>	<i>59</i>	<i>47</i>
2024	310	38	62	55	48	52	7	47	53	38

Table A4. 4. Sex composition by age and age composition of harvested sage-grouse within Wildlife Management Units with potential sage-grouse harvest, Oregon, 2024. Total by percent in each age class represents the overall proportion of the given age class in the harvest.

WMU ^a	Sample Size	Juveniles						Yearlings						Adults					
		Male		Female		Totals		Male		Female		Totals		Male		Female		Totals	
		N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
51 ^b	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
64 ^b	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
65	45	15	54	13	46	28	62	2	50	2	50	4	9	7	54	6	46	13	29
66	29	4	24	13	76	17	59	0	0	2	100	2	7	7	70	3	30	10	34
67	2	0	0	1	100	1	50	0	0	1	100	1	50	0	-	0	-	0	0
68A	33	7	47	8	53	15	45	2	67	1	33	3	9	5	33	10	67	15	45
68B	64	11	34	21	66	32	50	2	33	4	67	6	9	9	35	17	65	26	41
69	32	6	33	12	67	18	56	2	100	0	0	2	6	8	67	4	33	12	38
70	46	6	27	16	73	22	48	3	75	1	25	4	9	10	50	10	50	20	43
71 ^b	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
72	0	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-	0	-
73 ^c	11	4	57	3	43	7	64	0	-	0	-	0	0	2	50	2	50	4	36
74	48	12	41	17	59	29	60	0	0	1	100	1	2	7	39	11	61	18	38
<i>All</i>	<i>310</i>	<i>65</i>	<i>38</i>	<i>104</i>	<i>62</i>	<i>169</i>	<i>55</i>	<i>11</i>	<i>48</i>	<i>12</i>	<i>52</i>	<i>23</i>	<i>7</i>	<i>55</i>	<i>47</i>	<i>63</i>	<i>53</i>	<i>118</i>	<i>38</i>

^aWildlife Management Unit: 51 - Sumpter, 64 - Lookout Mtn., 65 - Beulah, 66 - Malheur River, 67 - Owyhee, 68A – Trout Creek Mtns, 68B – E. Whitehorse, 69 - Steens Mtn., 70 - Beatys Butte, 71 - Juniper, 72 - Silvies, 73 - Wagontire, 74 - Warner.

^bWMU no hunting permits offered in 2024.

^cWMU partially closed to hunting during 2024.

Table A4. 5. Sage-grouse production data as determined from hunter-harvested wings, Oregon, 1982–1984, 1989–1992, 1993–2023, and 2024. Data is divided into these periods due to the sage-grouse harvest closure during 1985–1988, and implementation of revised wing analysis methodology in 1993.

Year	n	% Juvenile	Chicks/Female	Chicks M:F
1982	73	53	2.4	26:74
1983	291	38	0.9	53:47
1984	144	40	1	42:58
1985–1988	Hunting Season Closed			
1989	326	41	1.1	46:54
1990	437	34	1	39:61
1991	295	31	0.8	37:63
1992	407	31	0.7	48:52
<i>1982–1992 Mean</i>	<i>282</i>	<i>38</i>	<i>1</i>	<i>48:58</i>
1993	439	47	1.4	51:49
1994	764	43	1.1	47:53
1995	456	36	0.8	42:58
1996	493	51	1.5	42:58
1997	586	54	1.8	47:53
1998	466	49	1.5	48:52
1999	671	56	2	46:54
2000	592	44	1.4	46:54
2001	670	54	1.9	50:50
2002	648	58	2.3	51:49
2003	655	48	1.6	46:54
2004	778	52	1.7	45:55
2005	829	45	1.4	46:54
2006	669	47	1.7	46:54
2007	485	28	0.6	44:56
2008	443	54	1.6	49:51
2009	493	57	2.3	47:53
2010	463	48	1.4	43:57
2011	422	53	1.3	43:57
2012	321	29	0.8	40:60
2013	254	58	2	50:50
2014	262	31	0.7	38:62
2015	290	58	2.3	43:57
2016	331	46	1.5	54:46
2017	270	39	1	42:58
2018	255	42	1.4	36:64
2019	145	48	1.4	44:56
2020	175	43	1.1	44:56

Table A4.5. Continued.

2021	150	41	1.1	46:54
2022	228	51	1.9	37:63
2023	247	61	2.5	57:43
<i>1993–2023 Mean</i>	<i>450</i>	<i>47</i>	<i>1.5</i>	<i>46:54</i>
2024	310	55	2.3	38:62

Nest Success and Production

Nest success was estimated based on wing molt patterns of adult and yearling females. Female sage-grouse replace primary feathers following completion of nesting activity (Braun and Schroeder 2015); thus, hens with successful nests will initiate their molt later than hens with unsuccessful nests. Wings from hens harvested while they were in the process of growing new primary feathers through P9 likely had a successful hatch (Braun and Schroeder 2015). Conversely, wings from hens which had unsuccessful nests are typically growing P10 or have completed their primary molt at the time of harvest because they began molting earlier (Braun and Schroeder 2015). In Oregon, P9 is used to estimate apparent nest success, except in years of extraordinary snow or drought. Use of P9 will generally give a minimum estimate of nest success, but in some years may underestimate actual nest success. Overall, apparent nest success in 2024 was 36%, lower than apparent nest success in 2023 (45%), and lower than the long-term average (42%; Table A4.6). The extent to which annual variability in apparent nest success is related to actual variability in nest success is unclear due to the impact of variable weather conditions and hatch dates, which may mask true changes in nest success. Apparent nest success in units where the number of adult or yearling hens was greater than 1 was highest in the Wagonfire WMU (100%, 2 of 2 total females), followed by the Beatys Butte MMU (64%, 7 of 11 females), and the Trout Creeks sub-unit of the Whitehorse WMU (55%, 6 of 11 total females; Table 7; Figure 2). Apparent nest success was low in several WMUs, including the Beulah WMU (0%; 0 of 8 females), Owyhee WMU (0%; 0 of 1 females), Warner WMU (17%; 2 of 12 females), and Malheur River WMU (20%; 1 of 5 females), decreasing the overall apparent nest success estimate (Figure 2). However, these estimates are likely driven by low sample sizes and may not be representative of the true nest success in these WMUs. The overall hatch appeared to be earlier in 2024, likely leading to more females initiating molt earlier than average. In this case, female retention of P10 may be a more appropriate indication of 2024 nest success, where

estimated nest success was 60% (Table A4.7). True nest success during 2024 was likely in the range of 36–60%.

Connelly et al. (2000) suggested that a chick per hen (CPH) ratio ≥ 2.25 indicates a healthy, stable or increasing population, but this ratio may be higher than required to maintain some populations and requires further study (Braun 2012). In Oregon, the long-term average (1993–2023) chick per hen ratio is 1.52 CPH. Production in 2024 as measured by chicks per hen was estimated at 2.25 CPH, well above the long-term average, and among highest CPH recorded in Oregon since hunter harvest sage-grouse wing data collection began in 1982 (Figure 3). Within WMUs, nest success was neither correlated with the proportion of juveniles in the harvest (Pearson's Correlation Coefficient = -0.05, $p = 0.90$) nor with the number of chicks per hen (Pearson's Correlation Coefficient = 0.03, $p = 0.93$). Historically, apparent nest success and production in Oregon have not been well correlated, with 2021 and 2023 serving as outliers, suggesting that nest success information derived from plumage characteristics is often unreliable. The pattern of disagreement between estimated P9 nest success and production as measured by the ratio of chicks per hen is historically consistent, with no significant correlation between the two values at the statewide level since current data collection began in 1993. The lack of correlation between these two values may suggest that variability in nesting and hatch dates due to climatological factors limits the utility of a single morphometric measure of nest success over time. Current methods for determining apparent nest success may be more useful as a measure of relative date of nesting between years than as a comparison of nest success among years. Further research is needed to develop accurate methods of determining nest success from wing data.

Table A4. 6. Sage-grouse nesting success as indicated by retention of at least primary feather P9, all wildlife management units with sage-grouse harvest, Oregon, 1993–2024.

Year	Nest Success (Retention of P9 and Lower)
1993	40
1994	40
1995	43
1996	51
1997	<i>No Data</i>
1998	30
1999	46
2000	45
2001	49
2002	47
2003	54
2004	35
2005	34
2006	49
2007	35
2008	48
2009	49
2010	37
2011	46
2012	63
2013	47
2014	52
2015	27
2016	30
2017	32
2018	46
2019	45
2020	33
2021	37
2022	39
2023	45
<i>31-Year Average</i>	<i>42</i>
2024	36

Table A4. 7. Sage-grouse nest success as indicated by retention of at least primary feather P9, and production rates, all WMUs with sage-grouse harvest, Oregon, 2024.

WMU	Adults			Yearling			All Hens			Harvest Age Composition			Production	
	Successful Adult Females (n)	Total Adult Females (n)	Adult Nest Success (%)	Successful Yearling Females (n)	Total Yearling Females (n)	Yearling Nest Success (%)	Successful Females (n)	Total Females (n)	Nest Success (%)	Total Harvest (n)	Juveniles in Harvest (n)	Juveniles in Harvest (%)	Juveniles per Female	Juveniles per Successful Female
Beatys Butte	6	10	60	1	1	100	7	11	64	46	22	48	2.00	3.14
Beulah	0	6	0	0	2	0	0	8	0	45	28	62	3.50	NA
E. Whitehorse	7	17	41	1	4	25	8	21	38	64	32	50	1.52	4.00
Malheur River	1	3	33	0	2	0	1	5	20	29	17	59	3.40	17.00
Owyhee	0	0	NA	0	1	0	0	1	0	2	1	50	1.00	NA
Silvies	0	0	NA	0	0	NA	0	0	NA	0	0	NA	NA	NA
Steens Mtn.	1	4	25	0	0	NA	1	4	25	32	18	56	4.50	18.00
Trout Creek Mtns.	5	10	50	1	1	100	6	11	55	33	15	45	1.36	2.50
Wagontire ^a	2	2	100	0	0	NA	2	2	100	11	7	64	3.50	3.50
Warner	2	11	18	0	1	0	2	12	17	48	29	60	2.42	14.50
All Areas (P9 or lower)	24	63	38	3	12	25	27	75	36	310	169	55	2.25	6.26
All Areas (P10 or lower)	35	63	56	10	12	83	45	75	60	310	169	55	2.25	3.76

^aWMU partially closed to hunting during 2024.




Figure A4. 3. Oregon sage-grouse production values (chicks per hen; CPH) and 32-year average (dashed line; 1993–2024) estimated from hunter harvested wing analyses, 1993–2024.

Hatching Chronology

Where possible, the length of the most recently replaced actively growing adult primary feather (usually P8 or P7) was recorded for juveniles. Ages of juveniles were calculated using growth data modified from captive-reared sage-grouse (Pyrah 1963). However, there is some evidence to suggest growth rates between wild and captive birds differ. Thus, the estimated hatch dates (Tables A4.8, A4.9) may be up to seven days earlier than the actual hatch date (Pyrah 1963).

Estimated chick hatch in 2024 began on 23 April, and lasted through late June, with two harvested juveniles hatching in July, on 1 and 9 July. Approximately 94% of males and 90% of females were estimated to have hatched prior to June 12 (Figure A4.5; Table A4.9). Typically, hatching chronology estimates in Oregon have suggested that peak male hatch occurs approximately 1 week prior to peak female hatch, likely due to flaws in the primary growth curves used to estimate hatch date. Hatching chronology estimates in 2024 indicate the first peak of male chick hatching occurred prior to May 1, with a second peak during June 5–11, while female chick hatching occurred during May 8–14 (Figure A4.4). Notably, 51% of males were estimated to have hatched prior to May 8, where only 27% of females were estimated to have hatched during this timeframe (Figure A4.5; Table A4.9). There is no known biological mechanism to explain sex-specific hatch dates, further supporting the need for improved sage-grouse chick growth curve data.

Table A4. 8. Estimated hatch dates for juvenile sage-grouse (% of total) from hunter-harvested wings, Oregon, 1993–2024.

Year	N	May					June				July		
		<1	1–7	8–14	15–21	22–28	29–4	5–11	12–18	19–25	26–2	3–9	10–16
1993	205	-	1	10	21	18	13	16	13	8	1	1	1
1994	327	1	9	22	18	16	11	12	8	3	1	1	-
1995	163	1	8	13	21	9	8	12	13	6	7	2	-
1996	253	2	9	15	12	14	11	17	10	4	2	3	1
1997	313	8	8	17	15	12	17	15	5	3	1	1	-
1998	229	2	10	13	15	18	14	10	3	7	2	4	1
1999	373	3	5	16	17	16	11	13	8	8	2	1	-
2000	260	7	7	17	18	16	15	14	4	2	1	-	-
2001	359	2	7	13	16	16	17	12	10	5	3	-	-
2002	373	5	6	17	13	21	13	13	4	4	3	1	-
2003	314	4	9	10	15	13	15	13	11	7	4	1	-
2004	398	3	10	24	24	14	11	8	5	2	1	-	-
2005	68	4	9	22	15	9	11	11	9	6	3	2	-
2006	323	1	3	10	12	12	18	21	15	7	1	1	-
2007	135	3	7	16	16	21	15	14	5	2	0	1	-
2008	241	3	7	10	12	15	15	15	8	8	7	-	-
2009	279	3	12	17	21	13	13	11	5	3	1	<1	-
2010	221	<2	6	9	18	13	15	14	13	4	4	<2	-
2011	178	-	<1	6	10	16	13	17	10	13	8	5	3
2012	94	3	5	25	14	16	11	10	11	5	-	-	-
2013	138	4	17	17	20	14	9	9	8	2	<1	-	-
2014	71	8	21	24	14	11	8	6	7	-	-	-	-
2015	152	3	17	28	15	9	14	8	6	1	1	-	-
2016	136	7	11	22	20	22	9	5	3	-	1	-	-
2017	96	6	6	17	21	20	9	11	6	2	1	-	-
2018	94	-	6	15	29	21	12	11	5	-	-	1	-
2019	67	15	6	12	13	16	10	15	7	3	1	-	-
2020	69	3	7	17	25	16	10	9	7	4	1	-	-
2021	57	3	7	24	11	5	3	3	1	-	-	-	-
2022	105	5	21	21	20	10	3	7	9	8	0	1	0
2023	140	16	15	19	18	18	12	24	9	5	1	3	0
2024	161	18	25	31	23	16	15	17	9	3	3	1	0

Table A4. 9. Estimated hatch dates, from hunter-harvested wings, for juvenile sage-grouse in Oregon, 2024.

Period	Males			Females			All Chicks		
	n	%	Cumulative %	n	%	Cumulative %	n	%	Cumulative %
< May 1	17	27	27	1	1	1	18	11	11
May 1–7	15	24	51	10	10	11	25	16	27
May 8–14	7	11	62	24	24	36	31	19	46
May 15–21	3	5	67	20	20	56	23	14	60
May 22–28	4	6	73	12	12	68	16	10	70
May 29–Jun 4	5	8	81	10	10	79	15	9	80
Jun 5–11	8	13	94	9	9	88	17	11	90
Jun 12–18	3	5	98	6	6	94	9	6	96
Jun 19–25	1	2	100	2	2	96	3	2	98
Jun 26–Jul 2	0	0	100	3	3	99	3	2	99
Jul 3–9	0	0	100	1	1	100	1	1	100
Jul 10–16	0	0	100	0	0	100	0	0	100




Figure A4. 4. Estimated hatch dates of male and female juvenile sage-grouse (% hatched during period), from hunter-harvested wings, Oregon, 2024.




Figure A4. 5. Estimated hatch dates of male and female juvenile sage-grouse (cumulative % hatched during period), from hunter-harvested wings, Oregon, 2024.

Annual Turnover

Analysis of annual turnover provides a rough estimate of adult mortality, assuming population stability. As in previous years, the data for annual turnover, when based on the proportion of yearlings in the fall harvest, were not able to support an unbiased analysis. Few yearlings were identifiable in 2024 ($n = 21$; 11 males, 12 females), likely due to early nesting in Oregon, corresponding to an early start of the primary molt. Generally, the timing of breeding and nesting in Oregon is earlier than those populations in the eastern portion of the sage-grouse distribution and in populations occupying higher elevations (Connelly et al. 2011), leading to a comparatively earlier molt in Oregon. Thus, the proportion of juveniles in the fall harvest of each sex was compared to the proportion of adults and yearlings (combined) of each sex to examine annual turnover (Table A4.10). This method is valid if one assumes the proportion of juveniles equals the annual loss of yearlings and adults. If the population was stable, annual mortality of adult and yearling males would be 51%, and annual mortality of adult and yearling females would be 44% (based on the 31-year average; Table A4.10).

Table A4. 10. Estimated annual turnover (%) of adult sage-grouse, assuming population stability, Oregon, 1993–2024.

Year	Males		Females	
	Young	Adults/Yearling	Young	Adults/Yearling
1993	54	46	41	59
1994	54	46	36	64
1995	44	56	31	69
1996	60	40	46	54
1997	61	39	49	51
1998	56	44	44	56
1999	60	40	52	48
2000	45	55	43	57
2001	61	39	49	51
2002	64	36	52	48
2003	50	50	47	53
2004	57	43	47	52
2005	47	53	43	57
2006	47	53	48	52
2007	33	67	25	75
2008	68	32	46	54
2009	58	42	55	45
2010	55	45	44	46
2011	42	58	43	57
2012	27	73	31	69
2013	69	31	50	50
2014	31	69	31	69
2015	61	39	56	44
2016	51	49	41	59
2017	45	55	36	64
2018	37	63	46	53
2019	54	46	44	56
2020	50	50	39	61
2021	48	52	37	63
2022	47	53	55	45
2023	70	30	53	48
<i>31-yr Avg</i>	<i>51</i>	<i>49</i>	<i>44</i>	<i>56</i>
2024	50	50	58	42

Conclusions

Oregon's sage-grouse hunting seasons are based on a long history of population monitoring and research. The current permit system allows ODFW to closely control legal harvest of sage-grouse. Each year, ODFW projects the fall population of sage-grouse based on lek counts. In 2024, ODFW estimated there were 19,624 sage-grouse in the fall population in the 9 WMUs where sage-grouse hunting is permitted (Table A4.11). ODFW offered 670 permits, of which 475 tags were purchased, and an estimated 436 individuals chose to hunt. The number of authorized permits increased for the 2024 hunting season compared to the 2022 and 2023 hunting seasons (640 permits) to reflect the population increase observed statewide in 2024. Additionally, production estimates were high in 2022 (1.92 CPH) and 2023 (2.50 CPH), supporting increased breeding and fall population estimates and leading to ODFW's increase in authorized permits for the 2024 hunting season. Despite the large increase in the estimated spring and fall population in the Beulah WMU, the number of permits was lowered (-25) in response to the Cow Valley and Durkee wildfires (July–August 2024), which significantly impacted sage-grouse habitat within the Beulah WMU.

In 2019, the methodology to estimate the fall sage-grouse population was updated to reflect the best available science. Historically, fall population estimates were generated using an estimate of chick production derived from summer brood route observations. However, analyses of brood route data revealed no long-term correlation between the chicks per hen ratios derived from brood routes and those derived from the wing analysis. Given this lack of correlation, brood route chick production estimates for sage-grouse in Oregon are likely unreliable and, thus, inappropriate to use during the harvest tag allocation process. In the place of brood route-derived estimates of chicks per hen, a conservative estimate of summer production (0.50 CPH) has been used to estimate fall population size in all units since 2019. The updated methodology lowered the estimated fall population size in all WMUs, and the number of permits offered in the Beatys Butte, Steens Mountain, and Warner WMUs were reduced in 2019. Permits in the other 5 WMUs were not reduced in 2019 due to harvest allocation in those units already existing well under the level necessary to maintain harvest at <5% of fall population size. This updated methodology has been used since 2019 to estimate the fall sage-grouse population and is likely to be used in the future.

ODFW has a self-imposed policy not to harvest more than 5% of the fall sage-grouse population, with harvest usually estimated at around 3% of the fall population. This harvest strategy is well within the guidelines suggested by the Western Association of Fish and Wildlife Agencies (Connelly et al. 2000). In addition, Oregon's harvest rate is well below the <11% harvest rate identified as unlikely to influence sage-grouse populations in Nevada and Colorado (Sedinger et al. 2010).

Compared to other states that offer a sage-grouse hunting season, Oregon's hunting season is likely the most conservative:

- Oregon's sage-grouse season is limited-entry for each WMU.
- Sage-grouse are not hunted range-wide in Oregon. Hunting is permitted in only 9 of 21 WMUs where sage-grouse occur (Figure A4.1).
- In certain cases, WMUs are divided into sub-units with separate permit allocations to address uneven distribution of hunters or other management or research needs (e.g. North/South Wagonfire sub-WMU and Whitehorse/Trout Creek Mountains sub-WMU).
- Permits are allocated to harvest no more than 5% of the estimated fall population in hunted WMUs (typically 3% or less in practice).
- Each permit holder is allowed to harvest only 2 sage-grouse per season.
- In 2024, estimated harvest of sage-grouse was 501 birds, 2.6% of the estimated 19,624 sage-grouse in hunted WMUs or sub-WMUs.

Through the collection of hunter-harvested wings, Oregon's sage-grouse hunting season provides crucial demographic data regarding the structure of sage-grouse populations in Oregon. This data would be costly or unfeasible to collect through other means.

Table A4. 11. Estimated fall sage-grouse population, maximum allowable harvest, hunter statistics, expected harvest, and permit allocation in Oregon wildlife management units (WMUs) where sage-grouse harvest is permitted, 2024.

WMU	Estimated Fall Population	Harvest Limit (5%)	Birds/Hunter ^a	Hunter Participation Rate ^a	2024 Permits	Expected Harvest (# birds)
Silvies	832	42	0.65	0.67	20	9
North Wagonfire	836	42	0.65	0.67	25	11
Beatys Butte	2,679	134	0.85	0.58	90	44
Steens Mtn.	1,534	77	0.85	0.58	50	25
Warner	1,928	96	0.85	0.58	70	34
Beulah	3,041	152	0.59	0.53	125	39
Malheur River	2,273	114	0.59	0.53	100	31
Owyhee	1,231	62	0.98	0.59	50	29
Trout Creek Mtn.	1,921	96	0.98	0.59	40	23
East Whitehorse	3,349	167	0.98	0.59	100	58
Total	19,624	982	-	-	670	303

^aHunter statistics based on average from hunter harvest survey by Data Analysis Unit (DAU) for years 2019–2023.

References

- Braun, C. E. 2012. Oregon sage-grouse wing analysis, 1993–2012. Unpubl. Report for Oregon Department of Fish and Wildlife. 40 p.
- Braun, C. E., D. A. Budeau, and M. A. Schroeder. 2015. Fall population structures of sage-grouse in Colorado and Oregon. Oregon Department of Fish and Wildlife Technical Report 005-2015. 72 p.
- Braun, C. E., and M. A. Schroeder. 2015. Age and sex identification from wings of hunter-harvested sage-grouse. *Wildlife Society Bulletin* 39:182–187.
- Connelly, J. W., C. A. Hagen, M. A. Schroeder. 2011. Characteristics and dynamics of greater sage-grouse populations. *Studies in Avian Biology* 38:53–67.
- Connelly, J. W., M. A. Schroeder, A. R. Sands, and C. E. Braun. 2000. Guidelines to manage sage grouse populations and their habitats. *Wildlife Society Bulletin* 28:967–985.
- Pyrah, D. B. 1963. Sage grouse investigations. Idaho Fish and Game Department. Wildlife Restoration Division Job Completion Report Project W-125-R-2. 71 p.
- Sedinger, J. S., G. C. White, S. Espinoza, E. T. Partee, and C. E. Braun. 2010. Assessing compensatory versus additive mortality: an example using greater sage-grouse. *Journal of Wildlife Management* 74:326–332.